Product Name: CD5 (10F9) Mouse Monoclonal Antibody Enkilife Catalog #: AMM00732

Summary

Production Name CD5 (10F9) Mouse Monoclonal Antibody

Description Primary antibody

Host Mouse
Application IHC-P

Reactivity Human, Rat, Mouse

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG1

Clonality Monoclonal Antibody

Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

Purification Affinity Purified

Immunogen

Gene Name CD5

CD5; LEU1; T-cell surface glycoprotein CD5; Lymphocyte antigen T1/Leu-1; CD antigen Alternative Names

CD5

Gene ID 921

SwissProt ID P06127

Application

Dilution Ratio IHC: 1/50-1/100

Molecular Weight -

Background




May act as a receptor in regulating T-cell proliferation. CD5 interacts with CD72/LYB-2.


Research Area

Immunology


Image Data



Immunohistochemistry analysis of paraffin-embedded Human livercancer tissue using CD5 (10F9) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody




Immunohistochemical analysis of paraffin-embedded Human tonsils using CD5 (10F9) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.




Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838



Immunohistochemistry analysis of paraffin-embedded mouse kidney tissue using CD5 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.



Immunofluorescence analysis of CD5 (10F9) in mouse heart tissue using CD5 (10F9) antibody(10G8)(red), and DAPI (blue).



Immunohistochemistry analysis of paraffin-embedded Human Tonsil Caricnoma using CD5 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Note

For research use only.