
**Monoclonal Antibody** Catalog #: AMM00752



# Summary

Acetyl-Histone H3 (Lys9) (10F10) Mouse Monoclonal Antibody **Production Name** 

Description Primary antibody

Host Mouse **Application** IHC-P

Reactivity Human, Rat, Mouse

### **Performance**

Conjugation Unconjugated Modification Acetylated

Isotype lgG1

**Clonality** Monoclonal Antibody

**Form** Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

**Buffer** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

**Purification Affinity Purified** 

## **Immunogen**

Gene Name H3C1

H3K9ac; H3/j; H3C1; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FJ; H3C10; H3C11;

**Alternative Names** HIST1H3J

8350

Gene ID SwissProt ID P68431


# **Application**

**Dilution Ratio** IHC: 1/50-1/100

**Molecular Weight** 

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

**Monoclonal Antibody** Catalog #: AMM00752




## **Background**

H3 Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability.


### Research Area

**Epigenetics and Nuclear Signaling** 

# **Image Data**



Immunohistochemistry analysis of paraffin-embedded Human Breast Carcinoma Tissue using Acetyl-Histone H3 (Lys9) (10F10) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunohistochemical analysis of paraffin-embedded Human tonsils using Acetyl-Histone H3 (Lys9) (10F10) antibody. Highpressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

## Note

For research use only.