

Summary

Production Name	ERK1 (1D10) Mouse Monoclonal Antibody
Description	Primary antibody
Host	Mouse
Application	IHC-P
Reactivity	Human, Rat, Mouse
Application	IHC-P

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG1
Clonality	Monoclonal Antibody
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Purification	Affinity Purified

Immunogen

Gene Name	MAPK3
Alternative Names	МАРК3
Gene ID	5595
SwissProt ID	P27361

Application

Dilution Ratio	IHC: 1/50-1/100
Molecular Weight	-

Background




Serine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway. MAPK1/ERK2 and MAPK3/ERK1 are the 2 MAPKs which play an important role in the MAPK/ERK cascade. They participate also in a signaling cascade initiated by activated KIT and KITLG/SCF. Depending on the cellular context, the MAPK/ERK cascade mediates diverse biological functions such as cell growth, adhesion, survival and differentiation through the regulation of transcription, translation, cytoskeletal rearrangements.


Research Area

Cell Biology

Image Data



Immunohistochemistry analysis of paraffin-embedded Human Tonsil Tissue using ERK1 (1D10) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunohistochemical analysis of paraffin-embedded Human tonsils using ERK1 (1D10) antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Note

For research use only.