

Summary

| Production Name | Cystatin C (6F12) Mouse Monoclonal Antibody |
|-----------------|---|
| Description | Primary antibody |
| Host | Mouse |
| Application | ELISA |
| Reactivity | Human |

Performance

| Conjugation | Unconjugated |
|--------------|--|
| Modification | Unmodified |
| lsotype | lgG1 |
| Clonality | Monoclonal Antibody |
| Form | Liquid |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw |
| | cycles. |
| Buffer | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3. |
| Purification | Affinity Purified |

Immunogen

| Gene Name | CST3 |
|-------------------|--|
| Alternative Names | CST3; Cystatin-C; Cystatin-3; Gamma-trace; Neuroendocrine basic polypeptide; Post- |
| | gamma-globulin |
| Gene ID | 1471 |
| SwissProt ID | P01034 |

Application

| Dilution Ratio | ELISA: 1/10000 |
|------------------|----------------|
| Molecular Weight | - |




Background

Cystatin C is a 14 kDa member of the Cystatin superfamily of cysteine protease inhibitors. Most cell types secrete Cystatin C. Cystatin C inhibits cathepsins, and thereby may function as a tumor suppressor by inhibiting cathepsin mediated tumor cell invasion. In addition, this tumor suppressor function can also be attributed to Cystatin C's ability to antagonize TGF-β1 signaling.

Research Area

Tags & Cell Markers

Image Data



Standard Curve for Human Cystatin C: Capture Antibody mouse mAb [6F12C7D8] to Human Cystatin C at 4ug/ml and Detector Antibody mouse mAb [7F6A5F3]to Human Cystatin C at 0.5ug/ml.

Note

For research use only.