

Summary

Production Name	PARP1 (7A1) Mouse Monoclonal Antibody	
Description	Primary antibody	
Host	Mouse	
Application	WB	
Reactivity	Human, Mouse, Rat, Chicken	

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG1
Clonality	Monoclonal Antibody
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Purification	Affinity Purified

Immunogen

Gene Name	PARP1
	PARP1; ADPRT; PPOL; Poly [ADP-ribose] polymerase 1; PARP-1; ADP-ribosyltransferase
Alternative Names	diphtheria toxin-like 1; ARTD1; NAD(+) ADP-ribosyltransferase 1; ADPRT 1; Poly[ADP-
	ribose] synthase 1
Gene ID	142
SwissProt ID	P09874

Application

Dilution Ratio	WB: 1/500-1/1000
Molecular Weight	Calculated MW: 113 kDa; Observed MW: 116 kDa




Background

Involved in the base excision repair (BER) pathway, by catalyzing the poly(ADP-ribosyl)ation of a limited number of acceptor proteins involved in chromatin architecture and in DNA metabolism. This modification follows DNA damages and appears as an obligatory step in a detection/signaling pathway leading to the reparation of DNA strand breaks.


Research Area

Epigenetics and Nuclear Signaling

Image Data



Western blot analysis of PARP in K562, Ramos, Jurkat lysates using PARP (7A1) antibody.



Western blot analysis of PARP1 (7A1) in CHO-K1, rat brain, C6, 3T3, Hela lysates using PARP (7A1) antibody.

Note

For research use only.