

Summary

Cystatin C (6A4) Mouse Monoclonal Antibody	
Primary antibody	
Mouse	
WB,IHC-P,ELISA	
Human	

Performance

Conjugation	Unconjugated	
Modification	Unmodified	
lsotype	lgG1	
Clonality	Monoclonal Antibody	
Form	Liquid	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw	
Storage	cycles.	
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.	
Purification	Affinity Purified	

Immunogen

Gene Name	CST3	
Alternative Names	CST3; Cystatin-C; Cystatin-3; Gamma-trace; Neuroendocrine basic polypeptide; Post-	
	gamma-globulin	
Gene ID	1471	
SwissProt ID	P01034	

Application

Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 ELISA: 1/10000
Molecular Weight	Calculated MW: 16 kDa; Observed MW: 16 kDa




Background

Cystatin C is a 14 kDa member of the Cystatin superfamily of cysteine protease inhibitors. Most cell types secrete Cystatin C. Cystatin C inhibits cathepsins, and thereby may function as a tumor suppressor by inhibiting cathepsin mediated tumor cell invasion. In addition, this tumor suppressor function can also be attributed to Cystatin C's ability to antagonize TGF-β1 signaling.


Research Area

Tags & Cell Markers

Image Data



Immunohistochemistry analysis of paraffin-embedded Human Brain Tissue using Cystatin C (6A4) antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Western blot analysis of Cystatin C (6A4) in NCIH460, SKOV-3, PC-3 lysates using Cystatin C (6A4) antibody

Note

For research use only.