

## Summary

Production Name	ERK1/2 (9A4) Mouse Monoclonal Antibody	
Description	Primary antibody	
Host	Mouse	
Application	WB,IHC-P	
Reactivity	Human, Rat, Mouse	

### Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG1
Clonality	Monoclonal Antibody
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Purification	Affinity Purified

### Immunogen

Gene Name	МАРКЗ/МАРК1	
	MAPK3; ERK1; ERT2; ERK-1; PRKM3; P44ERK1; P44MAPK; HS44KDAP; HUMKER1A; p44-	
Alternative Names	ERK1; p44-MAPK; MAPK1; ERK; p38; p40; p41; ERK2; ERT1; ERK-2; MAPK2; PRKM1;	
	PRKM2; P42MAPK; p41mapk; p42-MAPK.	
Gene ID	5595/5594	
SwissProt ID	P27361/P28482	

# Application

Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100
Molecular Weight	Calculated MW: 44,42 kDa; Observed MW: 44,42 kDa




#### Background

Serine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway. MAPK1/ERK2 and MAPK3/ERK1 are the 2 MAPKs which play an important role in the MAPK/ERK cascade. They participate also in a signaling cascade initiated by activated KIT and KITLG/SCF. Depending on the cellular context, the MAPK/ERK cascade mediates diverse biological functions such as cell growth, adhesion, survival and differentiation through the regulation of transcription, translation, cytoskeletal rearrangements.


#### **Research Area**

Cell Biology

## Image Data




Immunohistochemistry analysis of paraffin-embedded Human Breast Carcinoma using ERK1/2 (9A4) antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Western blot analysis of ERK1/2 (9A4) in Hela, C6 lysates using ERK1/2 (9A4) antibody





Western blot analysis of ERK1/2 (9A4) in mouse brain, rat brain , Hela lysates using ERK1/2 (9A4) antibody

#### Note

For research use only.