

# Summary

Production Name	GRP78 BiP (9E4) Mouse Monoclonal Antibody	
Description	Primary antibody	
Host	Mouse	
Application	WB,ICC/IF	
Reactivity	Human,Rat	

## Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG1
Clonality	Monoclonal Antibody
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Purification	Affinity Purified

## Immunogen

Gene Name	HSPA5
	HSPA5; GRP78; 78 kDa glucose-regulated protein; GRP-78; Endoplasmic reticulum
Alternative Names	lumenal Ca(2+)-binding protein grp78; Heat shock 70 kDa protein 5; Immunoglobulin
	heavy chain-binding protein; BiP
Gene ID	3309
SwissProt ID	P11021

# Application

Dilution Ratio	WB: 1/500-1/1000 IF: 1/50-1/200
Molecular Weight	Calculated MW: 72 kDa; Observed MW: 78 kDa




### Background

When Chinese hamster K12 cells are starved of glucose, the synthesis of several proteins, called glucose-regulated proteins (GRPs), is markedly increased. Hendershot et al. (1994) (PubMed 8020977) pointed out that one of these, GRP78 (HSPA5), also referred to as 'immunoglobulin heavy chain-binding protein' (BiP), is a member of the heat-shock protein-70 (HSP70) family and is involved in the folding and assembly of proteins in the endoplasmic reticulum (ER).


#### **Research Area**

Tags & Cell Markers

### **Image Data**



Western blot analysis of BiP/GRP78 (Cterminus) in Hela, C6, Lncap and MDA-MB-468 lysates using BiP/GRP78 (Cterminus)



Immunocytochemistry analysis of GRP78 BiP (9E4) in HeLa using BiP/GRP78 (Cterminus) antibody.

#### Note

For research use only.