Product Name: Lipoamide Dehydrogenase Rabbit

Monoclonal Antibody Catalog #: AMRe01457



Summary

Production Name Lipoamide Dehydrogenase Rabbit Monoclonal Antibody

Description Recombinant Rabbit Monoclonal antibody

Host Rabbit

Application WB,IHC-F,IHC-P,ICC/IF

Reactivity Human, Mouse, Rat

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG

Clonality Monoclonal Antibody

Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw $\bf Storage$

cycles.

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% **Buffer**

BSA

Purification Affinity Purified

Immunogen

Gene Name DLD

Diaphorase; Dihydrolipoamide dehydrogenase; DLDD; DLDH; GCSL; LAD; lipoamide

dehydrogenase; Lipoamide reductase; Lipoyl dehydrogenase; PHE3

 Gene ID
 1738

 SwissProt ID
 P09622

Application

Dilution Ratio WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200

Molecular Weight Calculated MW: 54 kDa; Observed MW: 54 kDa

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Lipoamide Dehydrogenase Rabbit

Monoclonal Antibody Catalog #: AMRe01457




Background

Lipoamide dehydrogenase is a component of the glycine cleavage system as well as of the alpha-ketoacid dehydrogenase complexes. Involved in the hyperactivation of spermatazoa during capacitation and in the spermatazoal acrosome reaction.


Research Area

Signal Transduction

Image Data



Immunocytochemistry analysis of Lipoamide Dehydrogenase (green) in Hela using Lipoamide Dehydrogenase antibody, and DAPI(blue).




Western blot analysis of Lipoamide Dehydrogenase in K562, rat Brain, C6, 3T3, Hela lysates using Lipoamide Dehydrogenase antibody.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Lipoamide Dehydrogenase Rabbit

Monoclonal Antibody Catalog #: AMRe01457





Immunohistochemistry analysis of paraffin-embedded Human breast cancer tissue using Lipoamide Dehydrogenase antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Note

For research use only.