Product Name: Annexin A1 Rabbit Monoclonal Antibody Enkilife Catalog #: AMRe01649

Summary

Production Name Annexin A1 Rabbit Monoclonal Antibody

Description Recombinant Rabbit Monoclonal antibody

Host Rabbit

Application WB,IHC-F,IHC-P,ICC/IF,IP

Reactivity Human, Mouse, Rat

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG

Clonality Monoclonal Antibody

Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% **Buffer**

BSA

Purification Affinity Purified

Immunogen

Gene Name ANXA1

ANXA1; ANX1; LPC1; Annexin A1; Annexin I; Annexin-1; Calpactin II; Calpactin-2; Alternative Names

Chromobindin-9; Lipocortin I; Phospholipase A2 inhibitory protein; p35

Gene ID 301

SwissProt ID P04083

Application

Dilution Ratio WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/20

Molecular Weight Calculated MW: 39 kDa; Observed MW: 39 kDa

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838


Background

Plays important roles in the innate immune response as effector of glucocorticoid-mediated responses and regulator of the inflammatory process. Has anti-inflammatory activity (PubMed:8425544). Plays a role in glucocorticoid-mediated down-regulation of the early phase of the inflammatory response. Promotes resolution of inflammation and wound healing (PubMed:25664854). Functions at least in part by activating the formyl peptide receptors and downstream signaling cascades (PubMed:15187149, PubMed:25664854). Promotes chemotaxis of granulocytes and monocytes via activation of the formyl peptide receptors (PubMed:15187149). Contributes to the adaptive immune response by enhancing signaling cascades that are triggered by T-cell activation, regulates differentiation and proliferation of activated T-cells (PubMed:17008549). Promotes the differentiation of T-cells into Th1 cells and negatively regulates differentiation into Th2 cells (PubMed:17008549). Has no effect on unstimulated T cells (PubMed:17008549). Promotes rearrangement of the actin cytoskeleton, cell polarization and cell migration (PubMed:15187149). Negatively regulates hormone exocytosis via activation of the formyl peptide receptors and reorganization of the actin cytoskeleton (PubMed:19625660). Has high affinity for Ca2+ and can bind up to eight Ca2+ ions . Displays Ca2+-dependent binding to phospholipid membranes (PubMed:2532504, PubMed:8557678). Plays a role in the formation of phagocytic cups and phagosomes. Plays a role in phagocytosis by mediating the Ca2+-dependent interaction between phagosomes and the actin cytoskeleton .

Research Area

Signal Transduction


Image Data




Immunocytochemistry analysis of Annexin A1 (green) in A549 using Annexin A1 antibody, and DAPI(blue).

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Western blot analysis of Annexin A1 in K562, C6, 3T3, Hela lysates using Annexin A1 antibody.



Immunohistochemistry analysis of paraffin-embedded Human tonsil using Annexin A1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Note

For research use only.