## **Product Name: ATG4A Rabbit Monoclonal Antibody**

Catalog #: AMRe01691



## **Summary**

**Production Name** ATG4A Rabbit Monoclonal Antibody

**Description** Recombinant Rabbit Monoclonal antibody

**Host** Rabbit

**Application** WB,IHC-P,IP

Reactivity Human

## **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

**Clonality** Monoclonal Antibody

Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% **Buffer** 

BSA

**Purification** Affinity Purified

## **Immunogen**

Gene Name ATG4A

ATG4A; APG4A; AUTL2; Cysteine protease ATG4A; AUT-like 2 cysteine endopeptidase;

Alternative Names Autophagin-2; Autophagy-related cysteine endopeptidase 2; Autophagy-related

protein 4 homolog A; hAPG4A

 Gene ID
 115201

 SwissProt ID
 Q8WYN0

## **Application**

**Dilution Ratio** WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/20

Molecular Weight Calculated MW: 45 kDa; Observed MW: 45 kDa

# **Product Name: ATG4A Rabbit Monoclonal Antibody**

Catalog #: AMRe01691




## **Background**

The cysteine protease Atg4 is pivotal to autophagosome membrane generation and regulation. Atg4 primes the Atg8 homologue for lipidation by cleaving its carboxy terminus and exposing its glycine residue for E1-like enzyme Atg7. The Atg8 homologue is transferred to the E2-like enzyme Atg3 before forming the Atg8-PE conjugate. During later stages of autophagy, Atg4 can reverse this lipidation event by cleaving PE, thereby recycling the Atg8 homologue.

### **Research Area**

**Cell Biology** 

## **Image Data**



Western blot analysis of ATG4A in Jurkat lysates using ATG4A antibody.

#### Note

For research use only.