

Summary

Production Name	Phospho-CBL (Ser669) Rabbit Monoclonal Antibody	
Description	Recombinant Rabbit Monoclonal antibody	
Host	Rabbit	
Application	WB	
Reactivity	Human	

Performance

Conjugation	Unconjugated
Modification	Phosphorylated
lsotype	IgG
Clonality	Monoclonal Antibody
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05%
	BSA
Purification	Affinity Purified

Immunogen

Gene Name	CBL
	CBL; CBL2; RNF55; E3 ubiquitin-protein ligase CBL; Casitas B-lineage lymphoma proto-
Alternative Names	oncogene; Proto-oncogene c-Cbl; RING finger protein 55; Signal transduction protein
	CBL
Gene ID	867
SwissProt ID	P22681

Application

Dilution Ratio	WB: 1/500-1/1000
Molecular Weight	Calculated MW: 100 kDa; Observed MW: 120 kDa




Background

This gene is a proto-oncogene that encodes a RING finger E3 ubiquitin ligase. The encoded protein is one of the enzymes required for targeting substrates for degradation by the proteasome. This protein mediates the transfer of ubiquitin from ubiquitin conjugating enzymes (E2) to specific substrates. This protein also contains an N-terminal phosphotyrosine binding domain that allows it to interact with numerous tyrosine-phosphorylated substrates and target them for proteasome degradation. As such it functions as a negative regulator of many signal transduction pathways. This gene has been found to be mutated or translocated in many cancers including acute myeloid leukaemia. Mutations in this gene are also the cause of Noonan syndrome-like disorder

Research Area

Cell Biology

Image Data



Western blot analysis of Phospho-CBL (Ser669) in K562 lysates using Phospho-CBL (Ser669) antibody.

Note

For research use only.