Catalog #: AMRe02149



Summary

Production Name	IKK gamma Rabbit Monoclonal Antibody
Description	Recombinant Rabbit Monoclonal antibody
Host	Rabbit
Application	WB,ICC/IF,IP
Reactivity	Human, Mouse, Rat

Performance

Conjugation	Unconjugated	
Modification	Unmodified	
lsotype	IgG	
Clonality	Monoclonal Antibody	
Form	Liquid	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw	
	cycles.	
Buffer	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05%	
	BSA	
Purification	Affinity Purified	

Immunogen

Gene Name	IKBKG
Alternative Names	IKBKG; FIP3; NEMO; NF-kappa-B essential modulator; NEMO; FIP-3; IkB kinase-
	associated protein 1; IKKAP1; Inhibitor of nuclear factor kappa-B kinase subunit
	gamma; I-kappa-B kinase subunit gamma; IKK-gamma; IKKG; IkB kinase subunit
	gamma; NF
Gene ID	8517
SwissProt ID	Q9Y6K9

Application

Dilution Ratio	WB: 1/500-1/1000 IF: 1/50-1/200 IP: 1/20
Molecular Weight	Calculated MW: 48 kDa; Observed MW: 48 kDa

Product Name: IKK gamma Rabbit Monoclonal Antibody Catalog #: AMRe02149

Background


The NF-κB/Rel transcription factors are present in the cytosol in an inactive state, complexed with the inhibitory IκB proteins. Most agents that activate NF-κB do so through a common pathway based on phosphorylation-induced, proteasome-mediated degradation of IκB.

EnkiLife


Research Area

Cell Biology


Image Data



Western blot analysis of IKK gamma in 3T3, PC-12 lysates using IKK gamma antibody.



Western blot analysis of IKK gamma in K562, Hela lysates using IKK gamma antibody.



Immunocytochemistry analysis of IKK gamma(green) in Hela using IKK gamma antibody, and DAPI(blue)

Note For research use only.