

Summary

Production Name	c-Met Rabbit Monoclonal Antibody	
Description	Recombinant Rabbit Monoclonal antibody	
Host	Rabbit	
Application	WB,IHC-F,IHC-P,ICC/IF	
Reactivity	Human, Mouse, Rat	

Performance

Conjugation	Unconjugated	
Modification	Unmodified	
lsotype	IgG	
Clonality	Monoclonal Antibody	
Form	Liquid	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw	
	cycles.	
Buffer	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05%	
	BSA	
Purification	Affinity Purified	

Immunogen

Gene Name	MET	
Alternative Names	MET; Hepatocyte growth factor receptor; HGF receptor; HGF/SF receptor; Proto-	
	oncogene c-Met; Scatter factor receptor; SF receptor; Tyrosine-protein kinase Met	
Gene ID	4233	
SwissProt ID	P08581	

Application

Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200
Molecular Weight	Calculated MW: 156 kDa; Observed MW: 170 kDa

Product Name: c-Met Rabbit Monoclonal Antibody Catalog #: AMRe02248




Background

The proto-oncogene MET product is the hepatocyte growth factor receptor and encodes tyrosine-kinase activity. The primary single chain precursor protein is post-translationally cleaved to produce the alpha and beta subunits, which are disulfide linked to form the mature receptor.


Research Area

Signal Transduction


Image Data



Western blot analysis of Met (c-Met) in K562, C6, 3T3 lysates using c-Met antibody.



Immunocytochemistry analysis of Met (c-Met)(green) in Hela using Met (c-Met) antibody, and DAPI(blue)



Product Name: c-Met Rabbit Monoclonal Antibody Catalog #: AMRe02248



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using Met (c-Met) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Note For research use only.