

**Product Name: beta Catenin Rabbit Monoclonal Antibody**  
**Catalog #: AMRe03746**

---

## Summary

<b>Production Name</b>	beta Catenin Rabbit Monoclonal Antibody
<b>Description</b>	Recombinant Rabbit Monoclonal antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC-P,IP
<b>Reactivity</b>	Human,Mouse,Rat,Hamster

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal Antibody
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
<b>Purification</b>	Affinity Purified

## Immunogen

<b>Gene Name</b>	CTNNB1
<b>Alternative Names</b>	CTNNB1; CTNNB; OK/SW-cl.35; Catenin beta-1; Beta-catenin
<b>Gene ID</b>	1499
<b>SwissProt ID</b>	P35222

## Application

<b>Dilution Ratio</b>	WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/20
<b>Molecular Weight</b>	Calculated MW: 85 kDa; Observed MW: 85 kDa

**Product Name: beta Catenin Rabbit Monoclonal Antibody**  
**Catalog #: AMRe03746**


## Background

Beta-catenin is an adherens junction protein. Adherens junctions (AJs; also called the zonula adherens) are critical for the establishment and maintenance of epithelial layers, such as those lining organ surfaces. AJs mediate adhesion between cells, communicate a signal that neighboring cells are present, and anchor the actin cytoskeleton. In serving these roles, AJs regulate normal cell growth and behavior.


## Research Area

Signal Transduction

## Image Data




Western blot analysis of beta Catenin in mouse brain lysates using beta Catenin antibody.



Western blot analysis of beta Catenin in C6, CHO-K1, HeLa lysates using beta Catenin antibody.

**Product Name: beta Catenin Rabbit Monoclonal Antibody**  
**Catalog #: AMRe03746**

---



Immunohistochemistry analysis of paraffin-embedded Human Cholangiocarcinoma using beta Catenin antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

**Note**

For research use only.