

## Summary

<b>Production Name</b>	BRD4 Rabbit Monoclonal Antibody
<b>Description</b>	Recombinant Rabbit Monoclonal antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC-P,ICC/IF,IP
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal Antibody
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Purification</b>	Affinity Chromatography

## Immunogen

<b>Gene Name</b>	BRD4
<b>Alternative Names</b>	Brd4; CAP; HUNK1; MCAP; Bromodomain containing 4; chromosome associated protein
<b>Gene ID</b>	23476
<b>SwissProt ID</b>	O60885

## Application

<b>Dilution Ratio</b>	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/50
<b>Molecular Weight</b>	Calculated MW: 152 kDa; Observed MW: 152 kDa

**Product Name: BRD4 Rabbit Monoclonal Antibody**  
**Catalog #: AMRe03895**




## Background

Bromodomain-containing protein 4 (BRD4) is a member of the bromodomains and extra terminal (BET) family of proteins, which also includes BRD2, BRD3, and BRDT. BET family proteins contain two tandem bromodomains and an extra terminal (ET) domain, and bind acetyl lysine residues. BRD4 is a chromatin-binding protein with a preference for Lys14 on histone H3 as well as Lys5 and Lys12 on histone H4. BRD4 chromatin binding occurs throughout the cell cycle, including condensed mitotic chromosomes, when the majority of genes are silenced.

## Research Area

Epigenetics and Nuclear Signaling

## Image Data



Western blot analysis of BRD4 in HeLa lysates using BRD4 antibody.

## Note

For research use only.