

Summary

Production Name	Ferritin Light Chain Rabbit Monoclonal Antibody	
Description	Recombinant Rabbit Monoclonal antibody	
Host	Rabbit	
Application	WB,FC	
Reactivity	Human, Mouse, Rat	

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Monoclonal Antibody
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide
	and 0.05% BSA.
Purification	Affinity Purified

Immunogen

Gene Name	FTL
Alternative Names	Ferritin L chain; Ferritin L subunit; Ferritin light chain; Ferritin light polypeptide; ferritin
	light polypeptide like 3; FTL; LFTD; NBIA 3;
Gene ID	2512
SwissProt ID	P02792

Application

Dilution Ratio	WB: 1/500-1/1000 FC: 1/50-1/100
Molecular Weight	Calculated MW:20 kDa;Observed MW: 20 kDa




Background

Stores iron in a soluble, non-toxic, readily available form. Important for iron homeostasis. Iron is taken up in the ferrous form and deposited as ferric hydroxides after oxidation. Also plays a role in delivery of iron to cells. Mediates iron uptake in capsule cells of the developing kidney.

Research Area

Neuroscience

Image Data



Western blot analysis of Ferritin Light Chain in HepG2, HeLa lysates using Ferritin Light Chain antibody.

Note

For research use only.