

**Product Name: Phospho-Chk1 (S280) (9M10) Rabbit
Monoclonal Antibody
Catalog #: AMRe05874**



Summary

Production Name	Phospho-Chk1 (S280) (9M10) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human

Performance

Conjugation	Unconjugated
Modification	Phospho Antibody
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	CHEK1
Alternative Names	Checkpoint kinase 1; Chk1; Chk1; rad27;
Gene ID	1111.0
SwissProt ID	O14757.

Application

Dilution Ratio	WB 1:500-1:2000
Molecular Weight	54kDa

Background

Serine/threonine-protein kinase which is required for checkpoint-mediated cell cycle arrest and activation of DNA repair in response to the presence of DNA damage or unreplicated DNA. May also negatively regulate cell cycle progression during unperturbed cell cycles. Serine/threonine-protein kinase which is required for checkpoint-mediated cell cycle arrest and activation of DNA repair in response to the presence of DNA damage or unreplicated DNA (PubMed:[11535615](http://www.uniprot.org/citations/11535615), PubMed:[12446774](http://www.uniprot.org/citations/12446774), PubMed:[12399544](http://www.uniprot.org/citations/12399544), PubMed:[14559997](http://www.uniprot.org/citations/14559997), PubMed:[14988723](http://www.uniprot.org/citations/14988723), PubMed:[15311285](http://www.uniprot.org/citations/15311285), PubMed:[15665856](http://www.uniprot.org/citations/15665856), PubMed:[15650047](http://www.uniprot.org/citations/15650047), PubMed:[32357935](http://www.uniprot.org/citations/32357935)). May also negatively regulate cell cycle progression during unperturbed cell cycles (PubMed:[11535615](http://www.uniprot.org/citations/11535615), PubMed:[12446774](http://www.uniprot.org/citations/12446774), PubMed:[12399544](http://www.uniprot.org/citations/12399544), PubMed:[14559997](http://www.uniprot.org/citations/14559997), PubMed:[14988723](http://www.uniprot.org/citations/14988723), PubMed:[15311285](http://www.uniprot.org/citations/15311285), PubMed:[15665856](http://www.uniprot.org/citations/15665856), PubMed:[15650047](http://www.uniprot.org/citations/15650047)). This regulation is achieved by a number of mechanisms that together help to preserve the integrity of the genome (PubMed:[11535615](http://www.uniprot.org/citations/11535615), PubMed:[12446774](http://www.uniprot.org/citations/12446774), PubMed:[12399544](http://www.uniprot.org/citations/12399544), PubMed:[14559997](http://www.uniprot.org/citations/14559997), PubMed:[14988723](http://www.uniprot.org/citations/14988723), PubMed:[15311285](http://www.uniprot.org/citations/15311285), PubMed:[15665856](http://www.uniprot.org/citations/15665856), PubMed:[15650047](http://www.uniprot.org/citations/15650047)). Recognizes the substrate consensus sequence [R-X-X- S/T] (PubMed:[11535615](http://www.uniprot.org/citations/11535615), PubMed:[12446774](http://www.uniprot.org/citations/12446774), PubMed:[12399544](http://www.uniprot.org/citations/12399544), PubMed:[14559997](http://www.uniprot.org/citations/14559997), PubMed:[14988723](http://www.uniprot.org/citations/14988723), PubMed:[15311285](http://www.uniprot.org/citations/15311285), PubMed:[15665856](http://www.uniprot.org/citations/15665856), PubMed:[15650047](http://www.uniprot.org/citations/15650047)).

**Product Name: Phospho-Chk1 (S280) (9M10) Rabbit
Monoclonal Antibody
Catalog #: AMRe05874**



<http://www.uniprot.org/citations/14559997> target="_blank">14559997, PubMed:14988723, PubMed:15311285, PubMed:15665856, PubMed:15650047). Binds to and phosphorylates CDC25A, CDC25B and CDC25C (PubMed:9278511, PubMed:12676583, PubMed:14681206, PubMed:12676925, PubMed:12759351, PubMed:19734889, PubMed:14559997). Phosphorylation of CDC25A at 'Ser-178' and 'Thr-507' and phosphorylation of CDC25C at 'Ser-216' creates binding sites for 14-3-3 proteins which inhibit CDC25A and CDC25C (PubMed:9278511). Phosphorylation of CDC25A at 'Ser-76', 'Ser-124', 'Ser-178', 'Ser-279' and 'Ser-293' promotes proteolysis of CDC25A (PubMed:9278511, PubMed:12676583, PubMed:14681206, PubMed:12676925, PubMed:12759351, PubMed:19734889). Phosphorylation of CDC25A at 'Ser-76' primes the protein for subsequent phosphorylation at 'Ser-79', 'Ser-82' and 'Ser-88' by NEK11, which is required for polyubiquitination and degradation of CDC25A (PubMed:9278511, PubMed:19734889, PubMed:20090422). Inhibition of CDC25 leads to increased inhibitory tyrosine phosphorylation of CDK-cyclin complexes and blocks cell cycle progression (PubMed:9278511). Also phosphorylates NEK6 (PubMed:18728393). Binds to and phosphorylates RAD51 at 'Thr-309', which promotes the release of RAD51 from BRCA2 and enhances the association of RAD51 with chromatin, thereby promoting DNA repair by homologous recombination (PubMed:15665856). Phosphorylates multiple sites within the C-terminus of TP53, which promotes activation of TP53 by acetylation and promotes cell cycle arrest and suppression of cellular proliferation (PubMed:10673501, PubMed:15659650, PubMed:15659650, PubMed:15659650).


**Product Name: Phospho-Chk1 (S280) (9M10) Rabbit
Monoclonal Antibody
Catalog #: AMRe05874**



href="http://www.uniprot.org/citations/16511572" target="_blank">16511572). Also promotes repair of DNA cross-links through phosphorylation of FANCE (PubMed:17296736). Binds to and phosphorylates TLK1 at 'Ser-743', which prevents the TLK1-dependent phosphorylation of the chromatin assembly factor ASF1A (PubMed:12660173, PubMed:12955071). This may enhance chromatin assembly both in the presence or absence of DNA damage (PubMed:12660173, PubMed:12955071). May also play a role in replication fork maintenance through regulation of PCNA (PubMed:18451105). May regulate the transcription of genes that regulate cell-cycle progression through the phosphorylation of histones (By similarity). Phosphorylates histone H3.1 (to form H3T11ph), which leads to epigenetic inhibition of a subset of genes (By similarity). May also phosphorylate RB1 to promote its interaction with the E2F family of transcription factors and subsequent cell cycle arrest (PubMed:17380128). Phosphorylates SPRTN, promoting SPRTN recruitment to chromatin (PubMed:31316063). Reduces replication stress and activates the G2/M checkpoint, by phosphorylating and inactivating PABIR1/FAM122A and promoting the serine/threonine-protein phosphatase 2A-mediated dephosphorylation and stabilization of WEE1 levels and activity (PubMed:33108758).

Research Area

Image Data



Western blot analysis of Phospho-Chk1 (S280) expression in 293T treated with Calyculin A cell lysate.

Note

**Product Name: Phospho-Chk1 (S280) (9M10) Rabbit
Monoclonal Antibody
Catalog #: AMRe05874**



For research use only.