

**Product Name: Phospho-NAK/TBK1 (S172) (10J4) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe05953**

---

## Summary

<b>Production Name</b>	Phospho-NAK/TBK1 (S172) (10J4) Rabbit Monoclonal Antibody
<b>Description</b>	Rabbit Monoclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,ELISA
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Phospho Antibody
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	TBK1 {ECO:0000303 PubMed:10581243, ECO:0000312 HGNC:HGNC:11584}
<b>Alternative Names</b>	FTDALS4; NAK; T2K; Tbk1;
<b>Gene ID</b>	29110.0
<b>SwissProt ID</b>	Q9UHD2.

## Application

<b>Dilution Ratio</b>	WB 1:500-1:2000
<b>Molecular Weight</b>	84kDa

**Product Name: Phospho-NAK/TBK1 (S172) (10J4) Rabbit  
Monoclonal Antibody  
Catalog #: AMRe05953**

---

## Background

Serine/threonine protein involved in the signaling cascade converging to the activation of the transcription factor NF-kappa-B. May function as an IKK kinase, playing an essential role in the transcription of a subset of TNF-alpha-induced genes. Also mediates production of RANTES/CCL5 and interferon-beta/IFNB1. Serine/threonine kinase that plays an essential role in regulating inflammatory responses to foreign agents (PubMed: [12692549](http://www.uniprot.org/citations/12692549), PubMed: [14703513](http://www.uniprot.org/citations/14703513), PubMed: [18583960](http://www.uniprot.org/citations/18583960), PubMed: [12702806](http://www.uniprot.org/citations/12702806), PubMed: [15367631](http://www.uniprot.org/citations/15367631), PubMed: [10581243](http://www.uniprot.org/citations/10581243), PubMed: [11839743](http://www.uniprot.org/citations/11839743), PubMed: [15485837](http://www.uniprot.org/citations/15485837), PubMed: [21138416](http://www.uniprot.org/citations/21138416), PubMed: [25636800](http://www.uniprot.org/citations/25636800), PubMed: [23453971](http://www.uniprot.org/citations/23453971), PubMed: [23453972](http://www.uniprot.org/citations/23453972), PubMed: [23746807](http://www.uniprot.org/citations/23746807), PubMed: [26611359](http://www.uniprot.org/citations/26611359), PubMed: [32404352](http://www.uniprot.org/citations/32404352)). Following activation of toll-like receptors by viral or bacterial components, associates with TRAF3 and TANK and phosphorylates interferon regulatory factors (IRFs) IRF3 and IRF7 as well as DDX3X (PubMed: [12692549](http://www.uniprot.org/citations/12692549), PubMed: [14703513](http://www.uniprot.org/citations/14703513), PubMed: [18583960](http://www.uniprot.org/citations/18583960), PubMed: [12702806](http://www.uniprot.org/citations/12702806), PubMed: [15367631](http://www.uniprot.org/citations/15367631), PubMed: [25636800](http://www.uniprot.org/citations/25636800)). This activity allows subsequent homodimerization and nuclear translocation of the IRFs leading to transcriptional activation of pro-inflammatory and antiviral genes including IFNA and IFNB (PubMed: [12702806](http://www.uniprot.org/citations/12702806), PubMed: [15367631](http://www.uniprot.org/citations/15367631), PubMed: [25636800](http://www.uniprot.org/citations/25636800), PubMed: [32972995](http://www.uniprot.org/citations/32972995)). In order to establish such an antiviral state, TBK1 form several different complexes whose composition depends on the type of cell and cellular stimuli (PubMed: [23453971](http://www.uniprot.org/citations/23453971), PubMed: [23453971](http://www.uniprot.org/citations/23453971)).

**Product Name: Phospho-NAK/TBK1 (S172) (10J4) Rabbit  
Monoclonal Antibody  
Catalog #: AMRe05953**


---

<http://www.uniprot.org/citations/23453972> target="\_blank">23453972</a>, PubMed:<a href="http://www.uniprot.org/citations/23746807" target="\_blank">23746807</a>). Plays a key role in IRF3 activation: acts by first phosphorylating innate adapter proteins MAVS, STING1 and TICAM1 on their pLxIS motif, leading to recruitment of IRF3, thereby licensing IRF3 for phosphorylation by TBK1 (PubMed:<a href="http://www.uniprot.org/citations/25636800" target="\_blank">25636800</a>, PubMed:<a href="http://www.uniprot.org/citations/30842653" target="\_blank">30842653</a>). Phosphorylated IRF3 dissociates from the adapter proteins, dimerizes, and then enters the nucleus to induce expression of interferons (PubMed:<a href="http://www.uniprot.org/citations/25636800" target="\_blank">25636800</a>). Thus, several scaffolding molecules including FADD, TRADD, MAVS, AZI2, TANK or TBKBP1/SINTBAD can be recruited to the TBK1- containing-complexes (PubMed:<a href="http://www.uniprot.org/citations/21931631" target="\_blank">21931631</a>). Under particular conditions, functions as a NF-kappa-B effector by phosphorylating NF-kappa-B inhibitor alpha/NFKBIA, IKBKB or RELA to translocate NF-Kappa-B to the nucleus (PubMed:<a href="http://www.uniprot.org/citations/10783893" target="\_blank">10783893</a>, PubMed:<a href="http://www.uniprot.org/citations/15489227" target="\_blank">15489227</a>). Restricts bacterial proliferation by phosphorylating the autophagy receptor OPTN/Optineurin on 'Ser-177', thus enhancing LC3 binding affinity and antibacterial autophagy (PubMed:<a href="http://www.uniprot.org/citations/21617041" target="\_blank">21617041</a>). Phosphorylates SMCR8 component of the C9orf72-SMCR8 complex, promoting autophagosome maturation (PubMed:<a href="http://www.uniprot.org/citations/27103069" target="\_blank">27103069</a>). Phosphorylates and activates AKT1 (PubMed:<a href="http://www.uniprot.org/citations/21464307" target="\_blank">21464307</a>). Seems to play a role in energy balance regulation by sustaining a state of chronic, low-grade inflammation in obesity, wich leads to a negative impact on insulin sensitivity (By similarity). Attenuates retroviral budding by phosphorylating the endosomal sorting complex required for transport-I (ESCRT-I) subunit VPS37C (PubMed:<a href="http://www.uniprot.org/citations/21270402" target="\_blank">21270402</a>). Phosphorylates Borna disease virus (BDV) P protein (PubMed:<a href="http://www.uniprot.org/citations/16155125" target="\_blank">16155125</a>). Plays an essential role in the TLR3- and IFN-dependent control of herpes virus HSV-1 and HSV-2 infections in the central nervous system (PubMed:<a href="http://www.uniprot.org/citations/22851595" target="\_blank">22851595</a>).

## Research Area

## Image Data

**Product Name: Phospho-NAK/TBK1 (S172) (10J4) Rabbit  
Monoclonal Antibody  
Catalog #: AMRe05953**



Western blot analysis of NAK/TBK1 (phospho S172) expression in (1) HeLa cell lysate; (2) HeLa cell treated with Calyculin A .

**Note**

For research use only.