

**Product Name: 5HT2C Receptor (7F8) Rabbit
Monoclonal Antibody
Catalog #: AMRe06339**

Summary

Production Name	5HT2C Receptor (7F8) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	HTR2C
Alternative Names	5-HT-1C; 5-HT-2C; 5-HT1C; 5-HT2C; 5-HTR2C; 5HT1C; 5HT2C; 5HTR2C; 5Hydroxytryptamine 2C receptor; Htr1c; HTR2C;
Gene ID	3358.0
SwissProt ID	P28335.

Application

Dilution Ratio	WB 1:500-1:2000
Molecular Weight	52kDa


**Product Name: 5HT2C Receptor (7F8) Rabbit
Monoclonal Antibody
Catalog #: AMRe06339**

Background

This is one of the several different receptors for 5-hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone, and a mitogen. This receptor mediates its action by association with G proteins that activate a phosphatidylinositol-calcium second messenger system. G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a receptor for various drugs and psychoactive substances, including ergot alkaloid derivatives, 1-2,5,-dimethoxy-4-iodophenyl-2-aminopropane (DOI) and lysergic acid diethylamide (LSD). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors. Beta-arrestin family members inhibit signaling via G proteins and mediate activation of alternative signaling pathways. Signaling activates a phosphatidylinositol-calcium second messenger system that modulates the activity of phosphatidylinositol 3-kinase and down-stream signaling cascades and promotes the release of Ca(2+) ions from intracellular stores. Regulates neuronal activity via the activation of short transient receptor potential calcium channels in the brain, and thereby modulates the activation of pro-opiomelanocortin neurons and the release of CRH that then regulates the release of corticosterone. Plays a role in the regulation of appetite and eating behavior, responses to angiogenic stimuli and stress. Plays a role in insulin sensitivity and glucose homeostasis.

Research Area

Image Data



Western blot analysis of 5HT2C Receptor expression in (1) SH-SY5Y cell lysate; (2) Mouse kidney lysate.

Note

For research use only.