

**Product Name: AP2S1 (16G15) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe06977**



## Summary

|                        |  |
|------------------------|--|
| <b>Production Name</b> | AP2S1 (16G15) Rabbit Monoclonal Antibody |
| <b>Description</b>     | Rabbit Monoclonal Antibody               |
| <b>Host</b>            | Rabbit                                   |
| <b>Application</b>     | WB                                       |
| <b>Reactivity</b>      | Human,Mouse,Rat                          |

## Performance

|                     |  |
|---------------------|--|
| <b>Conjugation</b>  | Unconjugated   |
| <b>Modification</b> | Unmodified   |
| <b>Isotype</b>      | IgG  |
| <b>Clonality</b>    | Monoclonal   |
| <b>Form</b>         | Liquid   |
| <b>Storage</b>      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.   |
| <b>Buffer</b>       | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |
| <b>Purification</b> | Affinity purification  |

## Immunogen

|                          |  |
|--------------------------|--|
| <b>Gene Name</b>         | AP2S1  |
| <b>Alternative Names</b> | AP17; AP17 delta; Ap2s1; CLAPS2; Sigma2 adaptin; |
| <b>Gene ID</b>           | 1175.0   |
| <b>SwissProt ID</b>      | P53680.  |

## Application

|                         |                 |
|-------------------------|-----------------|
| <b>Dilution Ratio</b>   | WB 1:500-1:2000 |
| <b>Molecular Weight</b> | 17kDa           |

**Product Name: AP2S1 (16G15) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe06977**




## Background

Component of the adaptor protein complex 2 (AP-2). Adaptor protein complexes function in protein Transport via Transport vesicles in different membrane traffic pathways. Adaptor protein complexes are vesicle coat components and appear to be involved in cargo selection and vesicle formation. Component of the adaptor protein complex 2 (AP-2). Adaptor protein complexes function in protein Transport via Transport vesicles in different membrane traffic pathways. Adaptor protein complexes are vesicle coat components and appear to be involved in cargo selection and vesicle formation. AP-2 is involved in clathrin-dependent endocytosis in which cargo proteins are incorporated into vesicles surrounded by clathrin (clathrin-coated vesicles, CCVs) which are destined for fusion with the early endosome. The clathrin lattice serves as a mechanical scaffold but is itself unable to bind directly to membrane components. Clathrin-associated adaptor protein (AP) complexes which can bind directly to both the clathrin lattice and to the lipid and protein components of membranes are considered to be the major clathrin adaptors contributing the CCV formation. AP-2 also serves as a cargo receptor to selectively sort the membrane proteins involved in receptor-mediated endocytosis. AP-2 seems to play a role in the recycling of synaptic vesicle membranes from the presynaptic surface. AP-2 recognizes Y-X-X-[FILMV] (Y-X-X-Phi) and [ED]-X-X-X-L-[LI] endocytosis signal motifs within the cytosolic tails of transmembrane cargo molecules. AP-2 may also play a role in maintaining normal post-endocytic trafficking through the ARF6-regulated, non-clathrin pathway. The AP-2 alpha and AP-2 sigma subunits are thought to contribute to the recognition of the [ED]-X-X-X-L-[LI] motif (By similarity). May also play a role in extracellular calcium homeostasis.

## Research Area

## Image Data



Western blot analysis of AP2S1 expression in HeLa cell lysate.

## Note

For research use only.

**Product Name: AP2S1 (16G15) Rabbit Monoclonal  
Antibody**  
**Catalog #: AMRe06977**

---

