Product Name: Caspase-1 (16E17) Rabbit Monoclonal



Catalog #: AMRe07956



Summary

Production Name Caspase-1 (16E17) Rabbit Monoclonal Antibody

Description Rabbit Monoclonal Antibody

HostRabbitApplicationWB,ELISAReactivityHuman,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type
Buffer	preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.
	Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name CASP1

Alternative Names CASP-1; ICE; IL-1 beta converting enzyme; IL-1BC; IL1BCE; caspase-1;

 Gene ID
 834.0

 SwissProt ID
 P29466.

Application

Dilution Ratio WB 1:500-1:2000

Molecular Weight 45kDa

Product Name: Caspase-1 (16E17) Rabbit Monoclonal

Antibody

Catalog #: AMRe07956



Background

Thiol protease that cleaves IL-1 beta between an Asp and an Ala, releasing the mature cytokine which is involved in a variety of inflammatory processes. Important for defense against pathogens. Cleaves and activates sterol regulatory element binding proteins (SREBPs). Can also promote apoptosis. Thiol protease involved in a variety of inflammatory processes by proteolytically cleaving other proteins, such as the precursors of the inflammatory cytokines interleukin-1 beta (IL1B) and interleukin 18 (IL18) as well as the pyroptosis inducer Gasdermin-D (GSDMD), into active mature peptides (PubMed: 15326478, PubMed: 1574116, PubMed:7876192, PubMed:15498465, PubMed:26375003, PubMed:32051255). Plays a key role in cell immunity as an inflammatory response initiator: once activated through formation of an inflammasome complex, it initiates a proinflammatory response through the cleavage of the two inflammatory cytokines IL1B and IL18, releasing the mature cytokines which are involved in a variety of inflammatory processes (PubMed: 1574116, PubMed:7876192, PubMed:15498465, PubMed:15326478, PubMed: 32051255). Cleaves a tetrapeptide after an Asp residue at position P1 (PubMed: 1574116 , PubMed:7876192, PubMed:15498465). Also initiates pyroptosis, a programmed lytic cell death pathway, through cleavage of GSDMD (PubMed: 26375003). In contrast to cleavage of interleukins IL1B and IL1B, recognition and cleavage of GSDMD is not strictly dependent on the consensus cleavage site but depends on an exosite interface on CASP1 that recognizes and binds the Gasdermin-D, C-terminal (GSDMD-CT) part (PubMed:32051255, PubMed:32109412, PubMed:32553275). Upon inflammasome activation, during DNA virus infection but not RNA virus challenge, controls antiviral immunity through the cleavage of CGAS, rendering it inactive (PubMed:28314590). In apoptotic cells, cleaves SPHK2 which is released from cells and remains enzymatically active extracellularly (PubMed: 20197547).

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Caspase-1 (16E17) Rabbit Monoclonal


Antibody

Catalog #: AMRe07956



Research Area

Image Data



Western blot analysis of Caspase-1 in HeLa cell lysate.

Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838