

# Summary

Production Name	Caspase-6 (13P13) Rabbit Monoclonal Antibody	
Description	Rabbit Monoclonal Antibody	
Host	Rabbit	
Application	WB	
Reactivity	Human,Mouse,Rat	

#### Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
Purification	Affinity purification

### Immunogen

Gene Name	CASP6
Alternative Names	MCH2; CASP6; Caspase-6;
Gene ID	839.0
SwissProt ID	P55212.A synthetic peptide of human Caspase-6

## Application

Dilution Ratio	WB: 1:1000
Molecular Weight	33kDa

## Background



Caspase-6 (Mch2) is one of the major executioner caspases functioning in cellular apoptotic processes. Upon apoptotic stimulation, initiator caspases such as caspase-9 are cleaved and activated. The activated upstream caspases further process downstream executioner caspases, such as caspase-3 and caspase-6, by cleaving them into large and small subunits, thereby initiating a caspase cascade leading to apoptosis. One of the major targets for caspase-6 is the membrane associated protein lamin A. Cysteine protease that plays essential roles in programmed cell death, axonal degeneration, development and innate immunity (PubMed:<a href="http://www.uniprot.org/citations/8663580" target=" blank">8663580</a>, PubMed:<a href="http://www.uniprot.org/citations/32298652" target=" blank">32298652</a>). During apoptosis, localizes in the nucleus and cleaves the nuclear structural protein NUMA1 and lamin A/LMNA thereby inducing nuclear shrinkage and fragmentation (PubMed:<a href="http://www.uniprot.org/citations/17401638" target=" blank">17401638</a>, PubMed:<a href="http://www.uniprot.org/citations/8663580" target=" blank">8663580</a>, PubMed:<a href="http://www.uniprot.org/citations/9463409" target=" blank">9463409</a>). Furthermore, cleaves many transcription factors such as NF-kappa-B and cAMP response element-binding protein/CREBBP (PubMed:<a href="http://www.uniprot.org/citations/10559921" target=" blank">10559921</a>, PubMed:<a href="http://www.uniprot.org/citations/14657026" target=" blank">14657026</a>). Cleaves phospholipid scramblase proteins XKR4 and XKR9 (By similarity). Plays an essential role in axon degeneration during axon pruning which is the remodeling of axons during neurogenesis but not apoptosis (By similarity). Regulates B-cell programs both during early development and after antigen stimulation (By similarity). In addition, promotes the ZBP1-mediated activation of programmed cell death pathways including pyroptosis, apoptosis, and necroptosis (PANoptosis) and plays an essential role in defense against viruses (PubMed: <a href="http://www.uniprot.org/citations/32298652" target=" blank">32298652</a>). Mechanistically, interacts with RIPK3 and enhances the interaction between RIPK3 and ZBP1, leading to ZBP1-mediated inflammasome activation and cell death (PubMed: <a href="http://www.uniprot.org/citations/32298652" target=" blank">32298652</a>).

# **Research Area**

#### Image Data





Western blot analysis of extracts from Jurkat cells using RM4582 at 1:1000.

#### Note

For research use only.