



Product Name: CDK9 (5L11) Rabbit Monoclonal Antibody
Catalog #: AMRe08574

Summary

Production Name	CDK9 (5L11) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
Purification	Affinity purification

Immunogen

Gene Name	CDK9
Alternative Names	Cyclin-dependent kinase 9; C-2K; Cell division cycle 2-like protein kinase 4; Cell division protein kinase 9; CDC2L4; TAK;
Gene ID	1025.0
SwissProt ID	P50750.A synthetic peptide of human Cdk9

Application

Dilution Ratio	WB: 1:1000
Molecular Weight	43kDa



Product Name: CDK9 (5L11) Rabbit Monoclonal Antibody
Catalog #: AMRe08574

Background

Cyclin dependent kinases (CDKs) are activated in part by cyclin binding and by phosphorylation of a conserved threonine in the T-loop domain. Member of the cyclin-dependent kinase pair (CDK9/cyclin-T) complex, also called positive transcription elongation factor b (P-TEFb), which facilitates the transition from abortive to production elongation by phosphorylating the CTD (C-terminal domain) of the large subunit of RNA polymerase II (RNAP II), SUPT5H and RDBP. Protein kinase involved in the regulation of transcription (PubMed:10574912, PubMed:10757782, PubMed:11145967, PubMed:11575923, PubMed:11809800, PubMed:11884399, PubMed:14701750, PubMed:16109376, PubMed:16109377, PubMed:20930849, PubMed:28426094). Member of the cyclin-dependent kinase pair (CDK9/cyclin-T) complex, also called positive transcription elongation factor b (P-TEFb), which facilitates the transition from abortive to productive elongation by phosphorylating the CTD (C-terminal domain) of the large subunit of RNA polymerase II (RNAP II) POLR2A, SUPT5H and RDBP (PubMed:10574912, PubMed:10757782, PubMed:11145967, PubMed:11575923, PubMed:11809800, PubMed:11884399, PubMed:14701750, PubMed:16109376, PubMed:16109377, PubMed:20930849, PubMed:28426094). This complex is inactive when in the 7SK snRNP complex form (PubMed:10574912, PubMed:10757782, PubMed:11145967, PubMed:11575923, PubMed:11809800, PubMed:11884399, PubMed:14701750, PubMed:16109376, PubMed:16109377, PubMed:20930849, PubMed:28426094).



Product Name: CDK9 (5L11) Rabbit Monoclonal Antibody
Catalog #: AMRe08574


href="http://www.uniprot.org/citations/11884399" target="_blank">>11884399, PubMed:>14701750, PubMed:>16109376, PubMed:>16109377, PubMed:>20930849, PubMed:>28426094). Phosphorylates EP300, MYOD1, RPB1/POLR2A and AR and the negative elongation factors DSIF and NELF (PubMed:>9857195, PubMed:>10912001, PubMed:>11112772, PubMed:>12037670, PubMed:>20081228, PubMed:>20980437, PubMed:>21127351). Regulates cytokine inducible transcription networks by facilitating promoter recognition of target transcription factors (e.g. TNF-inducible RELA/p65 activation and IL-6- inducible STAT3 signaling) (PubMed:>17956865, PubMed:>18362169). Promotes RNA synthesis in genetic programs for cell growth, differentiation and viral pathogenesis (PubMed:>10393184, PubMed:>11112772). P-TEFb is also involved in cotranscriptional histone modification, mRNA processing and mRNA export (PubMed:>15564463, PubMed:>19575011, PubMed:>19844166). Modulates a complex network of chromatin modifications including histone H2B monoubiquitination (H2Bub1), H3 lysine 4 trimethylation (H3K4me3) and H3K36me3; integrates phosphorylation during transcription with chromatin modifications to control co-transcriptional histone mRNA processing (PubMed:>15564463, PubMed:>19575011, PubMed:>19844166). The CDK9/cyclin-K complex has also a kinase activity towards CTD of RNAP II and can substitute for CDK9/cyclin-T P-TEFb in vitro (PubMed:>21127351). Replication stress response protein; the CDK9/cyclin-K complex is required for genome integrity maintenance, by promoting cell cycle recovery from replication arrest and limiting single-stranded DNA amount in response to replication stress, thus reducing the breakdown of stalled replication forks and avoiding DNA damage (PubMed:>20493174). In addition, probable function in DNA repair of isoform 2 via interaction with

Product Name: CDK9 (5L11) Rabbit Monoclonal Antibody
Catalog #: AMRe08574

KU70/XRCC6 (PubMed:[20493174](http://www.uniprot.org/citations/20493174)). Promotes cardiac myocyte enlargement (PubMed:[20081228](http://www.uniprot.org/citations/20081228)). RPB1/POLR2A phosphorylation on 'Ser-2' in CTD activates transcription (PubMed:[21127351](http://www.uniprot.org/citations/21127351)). AR phosphorylation modulates AR transcription factor promoter selectivity and cell growth. DSIF and NELF phosphorylation promotes transcription by inhibiting their negative effect (PubMed:[9857195](http://www.uniprot.org/citations/9857195), PubMed:[10912001](http://www.uniprot.org/citations/10912001), PubMed:[11112772](http://www.uniprot.org/citations/11112772)). The phosphorylation of MYOD1 enhances its transcriptional activity and thus promotes muscle differentiation (PubMed:[12037670](http://www.uniprot.org/citations/12037670)).

Research Area

Image Data



Western blot detection of Cdk9 in Hela,MCF-7,U2OS,U251 using Cdk9 antibody(1:1000 diluted)

Note

For research use only.