

Summary

Production Name	CEA(CD66e) (7M6) Rabbit Monoclonal Antibody	
Description	Rabbit Monoclonal Antibody	
Host	Rabbit	
Application	WB,ELISA	
Reactivity	Human	

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New typepreservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	CEACAM5
Alternative Names	adhesion molecule 5; CD66e; Ceacam5; Meconium antigen 100;
Gene ID	1048.0
SwissProt ID	P06731.

Application

Dilution Ratio	WB 1:500-1:2000
Molecular Weight	77kDa




Background

Carcinoembryonic antigen (CEA), also known as CD66e or CEACAM5, is a 180-200 kDa cell surface glycoprotein whose expression is elevated in intestinal carcinomas and other tumors. CEA mediates cell adhesion, though little more is known about its biological activity. Cell surface glycoprotein that plays a role in cell adhesion, intracellular signaling and tumor progression (PubMed:2803308, PubMed:10910050, PubMed:10910050, PubMed:10910050, PubMed:2803308, PubMed:10864933, PubMed:2803308, PubMed:2803308, PubMed:2803308, PubMed:2803308, PubMed

Research Area

Image Data



Western blot analysis of CEA(CD66e) expression in Human colon cancer lysate.

Note

For research use only.