

Summary

Production Name	COUP TF1 (17W3) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat
Host Application	Rabbit WB,ELISA

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New typepreservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	NR2F1
Alternative Names	COUP-TF1; EAR3; ERBAL3; NR2F1; NR2F2; Nuclear receptor subfamily 2 group F
Alternative Names	member 1; SVP44; TCFCOUP1; TFCOUP1;
Gene ID	7025.0
SwissProt ID	P10589.

Application

Dilution Ratio	WB 1:500-1:1000
Molecular Weight	46kDa




Background

Coup (chicken ovalbumin upstream promoter) transcription factor binds to the ovalbumin promoter and, in conjunction with another protein (S300-II) stimulates initiation of transcription. Binds to both direct repeats and palindromes of the 5'-AGGTCA-3' motif. Represses transcriptional activity of LHCG. Coup (chicken ovalbumin upstream promoter) transcription factor binds to the ovalbumin promoter and, in conjunction with another protein (S300-II) stimulates initiation of transcription. Binds to both direct repeats and palindromes of the 5'-AGGTCA-3' motif. Represses transcriptional activity of LHCG.

Research Area

Image Data



Western blot analysis of COUP TF1 expression in HEK293 cell lysate.

Note

For research use only.