

# Summary

| Production Name | Ctip1 (6S12) Rabbit Monoclonal Antibody |
|-----------------|---|
| Description     | Rabbit Monoclonal Antibody              |
| Host            | Rabbit                                  |
| Application     | WB,ELISA                                |
| Reactivity      | Human, Mouse, Rat                       |
|                 |   |

#### Performance

| Conjugation  | Unconjugated   |
|--------------|--|
| Modification | Unmodified   |
| lsotype      | IgG  |
| Clonality    | Monoclonal   |
| Form         | Liquid   |
| Storage      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.   |
| Buffer       | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New typepreservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.Avoid freeze / thaw cycle. |
| Purification | Affinity purification  |

# Immunogen

| Gene Name         | BCL11A   |
|-------------------|--|
| Alternative Names | EVI9; CTIP1; ZNF856; HBFQTL5; BCL11A-L; BCL11A-S; BCL11a-M; BCL11A-XL; |
| Gene ID           | 53335.0  |
| SwissProt ID      | Q9H165.  |

# Application

| Dilution Ratio   | WB 1:1000-1:5000 |
|------------------|------------------|
| Molecular Weight | 91kDa            |




#### Background

Functions as a myeloid and B-cell proto-oncogene. May play important roles in leukemogenesis and hematopoiesis. An essential factor in lymphopoiesis, is required for B-cell formation in fetal liver. May function as a modulator of the transcriptional repression activity of ARP1. Transcription factor (PubMed:<a href="http://www.uniprot.org/citations/16704730" target=" blank">16704730</a>, PubMed:<a href="http://www.uniprot.org/citations/29606353" target=" blank">29606353</a>). Associated with the BAF SWI/SNF chromatin remodeling complex (PubMed:<a href="http://www.uniprot.org/citations/23644491" target=" blank">23644491</a>). Binds to the 5'-TGACCA-3' sequence motif in regulatory regions of target genes, including a distal promoter of the HBG1 hemoglobin subunit gamma-1 gene (PubMed:<a href="http://www.uniprot.org/citations/29606353" target=" blank">29606353</a>). Involved in regulation of the developmental switch from gamma- to beta-globin, probably via direct repression of HBG1; hence indirectly repressing fetal hemoglobin (HbF) level (PubMed:<a href="http://www.uniprot.org/citations/29606353" target=" blank">29606353</a>, PubMed:<a href="http://www.uniprot.org/citations/26375765" target=" blank">26375765</a>). Involved in brain development (PubMed:<a href="http://www.uniprot.org/citations/27453576" target="\_blank">27453576</a>). May play a role in hematopoiesis (By similarity). Essential factor in lymphopoiesis required for B-cell formation in fetal liver (By similarity). May function as a modulator of the transcriptional repression activity of NR2F2 (By similarity).

## **Research Area**

## Image Data



Western blot analysis of Ctip1 expression in Jurkat cell lysate.

# Note

For research use only.

