

Product Name: CYP27A1 (5H15) Rabbit Monoclonal Antibody
Catalog #: AMRe09642



Summary

Production Name	CYP27A1 (5H15) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	CYP27A1 {ECO:0000303 PubMed:21411718, ECO:0000312 HGNC:HGNC:2605}
Alternative Names	CP27; CTX; CYP; CYP27; CYP27A1;
Gene ID	1593.0
SwissProt ID	Q02318.

Application

Dilution Ratio	WB 1:500-1:2000
Molecular Weight	60kDa

Product Name: CYP27A1 (5H15) Rabbit Monoclonal Antibody
Catalog #: AMRe09642




Background

Catalyzes the first step in the oxidation of the side chain of sterol intermediates; the 27-hydroxylation of 5-beta-cholestane-3-alpha,7-alpha,12-alpha-triol. Has also a vitamin D3-25-hydroxylase activity. Cytochrome P450 monooxygenase that catalyzes regio- and stereospecific hydroxylation of cholesterol and its derivatives. Hydroxylates (with R stereochemistry) the terminal methyl group of cholesterol side-chain in a three step reaction to yield at first a C26 alcohol, then a C26 aldehyde and finally a C26 acid (PubMed: [9660774](http://www.uniprot.org/citations/9660774), PubMed: [12077124](http://www.uniprot.org/citations/12077124), PubMed: [21411718](http://www.uniprot.org/citations/21411718), PubMed: [28190002](http://www.uniprot.org/citations/28190002)). Regulates cholesterol homeostasis by catalyzing the conversion of excess cholesterol to bile acids via both the 'neutral' (classic) and the 'acid' (alternative) pathways (PubMed: [9660774](http://www.uniprot.org/citations/9660774), PubMed: [1708392](http://www.uniprot.org/citations/1708392), PubMed: [11412116](http://www.uniprot.org/citations/11412116), PubMed: [2019602](http://www.uniprot.org/citations/2019602), PubMed: [7915755](http://www.uniprot.org/citations/7915755), PubMed: [9186905](http://www.uniprot.org/citations/9186905), PubMed: [9790667](http://www.uniprot.org/citations/9790667)). May also regulate cholesterol homeostasis via generation of active oxysterols, which act as ligands for NR1H2 and NR1H3 nuclear receptors, modulating the transcription of genes involved in lipid metabolism (PubMed: [9660774](http://www.uniprot.org/citations/9660774), PubMed: [12077124](http://www.uniprot.org/citations/12077124)). Plays a role in cholestanol metabolism in the cerebellum. Similarly to cholesterol, hydroxylates cholestanol and may facilitate sterol diffusion through the blood-brain barrier to the systemic circulation for further degradation (PubMed: [28190002](http://www.uniprot.org/citations/28190002)). Also hydroxylates retinal 7- ketocholesterol, a noxious oxysterol with pro-inflammatory and pro- apoptotic effects, and may play a role in its elimination from the retinal pigment epithelium (PubMed: [21411718](http://www.uniprot.org/citations/21411718)). May play a redundant role in vitamin D biosynthesis. Catalyzes 25-hydroxylation of vitamin D3 that is required for its conversion to a functionally active form (PubMed: [15465040](http://www.uniprot.org/citations/15465040)).

Research Area

Image Data

Product Name: CYP27A1 (5H15) Rabbit Monoclonal Antibody
Catalog #: AMRe09642



Western blot analysis of CYP27A1 expression in HepG2 cell lysate.

Note

For research use only.