

Summary

Cytokeratin 8 (3B5) Rabbit Monoclonal Antibody	
Rabbit Monoclonal Antibody	
Rabbit	
WB,ELISA	
Human, Mouse	

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New typepreservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	KRT8	
Alternative Names	CARD2; CK-8; CK8; CYK8; cytokeratin 8; Cytokeratin-8; K2C8; K8; keratin 8; Keratin, type	
Alternative Names	II cytoskeletal 8; Keratin-8; KO; KRT8; Type-II keratin Kb8;	
Gene ID	3856.0	
SwissProt ID	P05787.	

Application

Dilution Ratio	WB 1:10000~1:20000
Molecular Weight	54kDa




Background

Together with KRT19, helps to link the contractile apparatus to dystrophin at the costameres of striated muscle. K8 a type II cytoskeletal keratin. The keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells and are subdivided into cytokeratins and hair keratins. Together with KRT19, helps to link the contractile apparatus to dystrophin at the costameres of striated muscle.

Research Area

Image Data



Western blot analysis of Cytokeratin 8 expression in HaCaT cell lysate.

Note

For research use only.