

Product Name: DKC1 (12X18) Rabbit Monoclonal Antibody
Catalog #: AMRe10001

Summary

| | |
|------------------------|---|
| Production Name | DKC1 (12X18) Rabbit Monoclonal Antibody |
| Description | Rabbit Monoclonal Antibody |
| Host | Rabbit |
| Application | WB |
| Reactivity | Human,Mouse,Rat |

Performance

| | |
|---------------------|--|
| Conjugation | Unconjugated |
| Modification | Unmodified |
| Isotype | IgG |
| Clonality | Monoclonal |
| Form | Liquid |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |
| Buffer | Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA. |
| Purification | Affinity purification |

Immunogen

| | |
|--------------------------|--|
| Gene Name | DKC1 |
| Alternative Names | DKC1; DKC; CBF5; DKCX; NAP57; NOLA4; XAP101; |
| Gene ID | 1736.0 |
| SwissProt ID | O60832.A synthetic peptide of human DKC1 |

Application

| | |
|-------------------------|-------------------|
| Dilution Ratio | WB: 1:1000-1:5000 |
| Molecular Weight | 58kDa |


Background

Product Name: DKC1 (12X18) Rabbit Monoclonal Antibody
Catalog #: AMRe10001

Required for ribosome biogenesis and telomere maintenance. Probable catalytic subunit of H/ACA small nucleolar ribonucleoprotein (H/ACA snoRNP) complex, which catalyzes pseudouridylation of rRNA. This involves the isomerization of uridine such that the ribose is subsequently attached to C5, instead of the normal N1. [Isoform 1]: Catalytic subunit of H/ACA small nucleolar ribonucleoprotein (H/ACA snoRNP) complex, which catalyzes pseudouridylation of rRNA (PubMed:25219674). This involves the isomerization of uridine such that the ribose is subsequently attached to C5, instead of the normal N1 (PubMed:25219674). Each rRNA can contain up to 100 pseudouridine ('psi') residues, which may serve to stabilize the conformation of rRNAs. Required for ribosome biogenesis and telomere maintenance (PubMed:19179534, PubMed:25219674). Also required for correct processing or intranuclear trafficking of TERC, the RNA component of the telomerase reverse transcriptase (TERT) holoenzyme (PubMed:19179534).

Research Area

Image Data



Western blot detection of DKC1 in K562, C6, 3T3, HeLa cell lysates using DKC1 antibody (1:1000 diluted).

Note

For research use only.