Product Name: Emi1 (16Z12) Rabbit Monoclonal

Antibody

Catalog #: AMRe10439



Summary

Production Name Emi1 (16Z12) Rabbit Monoclonal Antibody

Description Rabbit Monoclonal Antibody

Host Rabbit

Application WB,ELISA,IHC **Reactivity** Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type
Buffer	preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.
	Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name FBXO5

Alternative Names EMI1; FBX5; Fbxo31; fbxo5;

 Gene ID
 26271.0

 SwissProt ID
 Q9UKT4.

Application

Dilution Ratio WB 1:500-1:2000

Molecular Weight 50kDa

 Product Name: Emi1 (16Z12) Rabbit Monoclonal

Antibody

Catalog #: AMRe10439



Background

Regulates progression through early mitosis by inhibiting the anaphase promoting complex/cyclosome (APC). Binds to the APC activators CDC20 and FZR1/CDH1 to prevent APC activation. Can also bind directly to the APC to inhibit substratebinding. Regulator of APC activity during mitotic and meiotic cell cycle (PubMed: 17485488, PubMed:17234884, PubMed:17875940, PubMed:23708001, PubMed: 23708605, PubMed:16921029). During mitotic cell cycle plays a role as both substrate and inhibitor of APC-FZR1 complex (PubMed: 29875408, PubMed:17485488, PubMed:17234884, PubMed:17875940, PubMed:23708001, PubMed:23708605, PubMed:16921029). During G1 phase, plays a role as substrate of APC-FZR1 complex E3 ligase (PubMed:29875408). Then switches as an inhibitor of APC-FZR1 complex during S and G2 leading to cell-cycle commitment (PubMed:29875408). As APC inhibitor, prevents the degradation of APC substrates at multiple levels: by interacting with APC and blocking access of APC substrates to the Dbox coreceptor, formed by FZR1 and ANAPC10; by suppressing ubiquitin ligation and chain elongation by APC by preventing the UBE2C and UBE2S activities (PubMed: 23708605, PubMed:23708001, PubMed:16921029). Plays a role in genome integrity preservation by coordinating DNA replication with mitosis through APC inhibition in interphase to stabilize CCNA2 and GMNN in order to promote mitosis and prevent rereplication and DNA damage-induced cellular senescence (PubMed:17234884, PubMed:17485488, PubMed:17875940). During oocyte maturation, plays a role in meiosis through inactivation of APC-FZR1 complex. Inhibits APC through RPS6KA2 interaction that increases FBXO5 affiniy for CDC20 leading to the metaphase arrest of the second meiotic division before fertilization (By similarity). Controls entry into the first meiotic division through inactivation of APC-FZR1 complex (By similarity). Promotes migration and

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Emi1 (16Z12) Rabbit Monoclonal

Antibody


Catalog #: AMRe10439



osteogenic differentiation of mesenchymal stem cells (PubMed:29850565).

Research Area

Image Data



Western blot analysis of Emi1 expression in HepG2 cell lysate.

Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838