

**Product Name: Frizzled 8 (11I2) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe11137**



## Summary

|                        |  |
|------------------------|--|
| <b>Production Name</b> | Frizzled 8 (11I2) Rabbit Monoclonal Antibody |
| <b>Description</b>     | Rabbit Monoclonal Antibody                   |
| <b>Host</b>            | Rabbit                                       |
| <b>Application</b>     | WB,ELISA                                     |
| <b>Reactivity</b>      | Human,Mouse,Rat                              |

## Performance

|                     |  |
|---------------------|--|
| <b>Conjugation</b>  | Unconjugated   |
| <b>Modification</b> | Unmodified   |
| <b>Isotype</b>      | IgG  |
| <b>Clonality</b>    | Monoclonal   |
| <b>Form</b>         | Liquid   |
| <b>Storage</b>      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.   |
| <b>Buffer</b>       | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |
| <b>Purification</b> | Affinity purification  |

## Immunogen

|                          |                              |
|--------------------------|------------------------------|
| <b>Gene Name</b>         | FZD8                         |
| <b>Alternative Names</b> | Frizzled-8; FZ8; FZD8; hFz8; |
| <b>Gene ID</b>           | 8325.0                       |
| <b>SwissProt ID</b>      | Q9H461.                      |

## Application

|                         |                 |
|-------------------------|-----------------|
| <b>Dilution Ratio</b>   | WB 1:500-1:2000 |
| <b>Molecular Weight</b> | 73kDa           |

**Product Name: Frizzled 8 (11I2) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe11137**




## Background

Receptor for Wnt proteins. Component of the Wnt-Fzd-LRP5-LRP6 complex that triggers beta-catenin signaling through inducing aggregation of receptor-ligand complexes into ribosome-sized signalosomes. Receptor for Wnt proteins. Component of the Wnt-Fzd-LRP5-LRP6 complex that triggers beta-catenin signaling through inducing aggregation of receptor-ligand complexes into ribosome-sized signalosomes. The beta-catenin canonical signaling pathway leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues. Coreceptor along with RYK of Wnt proteins, such as WNT1.

## Research Area

## Image Data



Western blot analysis of Frizzled 8 expression in Jurkat cell lysate.

## Note

For research use only.