

Product Name: GluR2 (5C16) Rabbit Monoclonal Antibody
Catalog #: AMRe11492

Summary

Production Name	GluR2 (5C16) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
Purification	Affinity purification

Immunogen

Gene Name	GRIA2
Alternative Names	AMPA 2; AMPA selective glutamate receptor 2; AMPA2; GluA2; GLUR B; GluR K2; GLUR2; GLURB; Gria2; HBGR2;
Gene ID	2891.0
SwissProt ID	P42262.A synthetic peptide of human Ionotropic Glutamate receptor 2

Application

Dilution Ratio	WB: 1:1000-1:10000
Molecular Weight	99kDa


Product Name: GluR2 (5C16) Rabbit Monoclonal Antibody
Catalog #: AMRe11492

Background

Ionotropic glutamate receptor. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist. Receptor for glutamate that functions as ligand-gated ion channel in the central nervous system (PubMed:31300657). It plays an important role in excitatory synaptic transmission. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist. In the presence of CACNG4 or CACNG7 or CACNG8, shows resensitization which is characterized by a delayed accumulation of current flux upon continued application of glutamate. Through complex formation with NSG1, GRIP1 and STX12 controls the intracellular fate of AMPAR and the endosomal sorting of the GRIA2 subunit toward recycling and membrane targeting (By similarity).

Research Area

Image Data



Western blot detection of Ionotropic Glutamate receptor 2 in C6 cell lysates using Ionotropic Glutamate receptor 2 antibody(1:1000 diluted).

Note

For research use only.