

**Product Name: HAUSP / USP7 (17N8) Rabbit  
Monoclonal Antibody  
Catalog #: AMRe11905**

---

## Summary

<b>Production Name</b>	HAUSP / USP7 (17N8) Rabbit Monoclonal Antibody
<b>Description</b>	Rabbit Monoclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	USP7
<b>Alternative Names</b>	TEF1; HAUSP; USP7;
<b>Gene ID</b>	7874.0
<b>SwissProt ID</b>	Q93009.A synthetic peptide of human USP7

## Application

<b>Dilution Ratio</b>	WB: 1:1000
<b>Molecular Weight</b>	128kDa

## Background

---

**Product Name: HAUSP / USP7 (17N8) Rabbit  
Monoclonal Antibody  
Catalog #: AMRe11905**

---


Hydrolase that deubiquitinates target proteins such as FOXO4, p53/TP53, MDM2, ERCC6, DNMT1, UHRF1, PTEN and DAXX (PubMed:11923872, PubMed:15053880, PubMed:16964248, PubMed:18716620, PubMed:25283148). Hydrolase that deubiquitinates target proteins such as FOXO4, p53/TP53, MDM2, ERCC6, DNMT1, UHRF1, PTEN, KMT2E/MLL5 and DAXX (PubMed:<a href="http://www.uniprot.org/citations/11923872" target="\_blank">11923872</a>, PubMed:<a href="http://www.uniprot.org/citations/15053880" target="\_blank">15053880</a>, PubMed:<a href="http://www.uniprot.org/citations/16964248" target="\_blank">16964248</a>, PubMed:<a href="http://www.uniprot.org/citations/18716620" target="\_blank">18716620</a>, PubMed:<a href="http://www.uniprot.org/citations/25283148" target="\_blank">25283148</a>, PubMed:<a href="http://www.uniprot.org/citations/26678539" target="\_blank">26678539</a>, PubMed:<a href="http://www.uniprot.org/citations/28655758" target="\_blank">28655758</a>). Together with DAXX, prevents MDM2 self-ubiquitination and enhances the E3 ligase activity of MDM2 towards p53/TP53, thereby promoting p53/TP53 ubiquitination and proteasomal degradation (PubMed:<a href="http://www.uniprot.org/citations/15053880" target="\_blank">15053880</a>, PubMed:<a href="http://www.uniprot.org/citations/16845383" target="\_blank">16845383</a>, PubMed:<a href="http://www.uniprot.org/citations/18566590" target="\_blank">18566590</a>, PubMed:<a href="http://www.uniprot.org/citations/20153724" target="\_blank">20153724</a>). Deubiquitinates p53/TP53, preventing degradation of p53/TP53, and enhances p53/TP53-dependent transcription regulation, cell growth repression and apoptosis (PubMed:<a href="http://www.uniprot.org/citations/25283148" target="\_blank">25283148</a>). Deubiquitinates p53/TP53 and MDM2 and strongly stabilizes p53/TP53 even in the presence of excess MDM2, and also induces p53/TP53-dependent cell growth repression and apoptosis (PubMed:<a href="http://www.uniprot.org/citations/11923872" target="\_blank">11923872</a>). Deubiquitination of FOXO4 in presence of hydrogen peroxide is not dependent on p53/TP53 and inhibits FOXO4-induced transcriptional activity (PubMed:<a href="http://www.uniprot.org/citations/16964248" target="\_blank">16964248</a>). In association with DAXX, is involved in the deubiquitination and translocation of PTEN from the nucleus to the cytoplasm, both processes that are counteracted by PML (PubMed:<a href="http://www.uniprot.org/citations/18716620" target="\_blank">18716620</a>). Deubiquitinates KMT2E/MLL5 preventing KMT2E/MLL5 proteasomal-mediated degradation (PubMed:<a href="http://www.uniprot.org/citations/26678539" target="\_blank">26678539</a>). Involved in cell proliferation during early embryonic development. Involved in transcription-coupled nucleotide excision repair (TC-NER) in response to UV damage: recruited to DNA damage sites following interaction with KIAA1530/UVSSA and promotes deubiquitination of ERCC6, preventing UV- induced degradation of ERCC6 (PubMed:<a href="http://www.uniprot.org/citations/22466611" target="\_blank">22466611</a>, PubMed:<a href="http://www.uniprot.org/citations/22466612" target="\_blank">22466612</a>). Involved in maintenance of DNA methylation via its interaction with UHRF1 and DNMT1: acts by mediating deubiquitination of UHRF1 and DNMT1, preventing their degradation and promoting DNA methylation by DNMT1 (PubMed:<a href="http://www.uniprot.org/citations/21745816" target="\_blank">21745816</a>, PubMed:<a href="http://www.uniprot.org/citations/22411829" target="\_blank">22411829</a>). Deubiquitinates alkylation repair

**Product Name: HAUSP / USP7 (17N8) Rabbit  
Monoclonal Antibody  
Catalog #: AMRe11905**

enzyme ALKBH3. OTUD4 recruits USP7 and USP9X to stabilize ALKBH3, thereby promoting the repair of alkylated DNA lesions (PubMed:<a href="http://www.uniprot.org/citations/25944111" target="\_blank">25944111</a>). Acts as a chromatin regulator via its association with the Polycomb group (PcG) multiprotein PRC1-like complex; may act by deubiquitinating components of the PRC1-like complex (PubMed:<a href="http://www.uniprot.org/citations/20601937" target="\_blank">20601937</a>). Able to mediate deubiquitination of histone H2B; it is however unsure whether this activity takes place in vivo (PubMed:<a href="http://www.uniprot.org/citations/20601937" target="\_blank">20601937</a>). Exhibits a preference towards 'Lys-48'-linked ubiquitin chains (PubMed:<a href="http://www.uniprot.org/citations/22689415" target="\_blank">22689415</a>). Increases regulatory T-cells (Treg) suppressive capacity by deubiquitinating and stabilizing the transcription factor FOXP3 which is crucial for Treg cell function (PubMed:<a href="http://www.uniprot.org/citations/23973222" target="\_blank">23973222</a>). Plays a role in the maintenance of the circadian clock periodicity via deubiquitination and stabilization of the CRY1 and CRY2 proteins (PubMed:<a href="http://www.uniprot.org/citations/27123980" target="\_blank">27123980</a>). Deubiquitinates REST, thereby stabilizing REST and promoting the maintenance of neural progenitor cells (PubMed:<a href="http://www.uniprot.org/citations/21258371" target="\_blank">21258371</a>). Deubiquitinates SIRT7, inhibiting SIRT7 histone deacetylase activity and regulating gluconeogenesis (PubMed:<a href="http://www.uniprot.org/citations/28655758" target="\_blank">28655758</a>).

## Research Area

## Image Data



Western blot detection of USP7 in C6 cell lysates using USP7 antibody(1:1000 diluted).

## Note

For research use only.