# **Product Name: Hepcidin (16X4) Rabbit Monoclonal**

**Antibody** 

Catalog #: AMRe11984



# **Summary**

**Production Name** Hepcidin (16X4) Rabbit Monoclonal Antibody

**Description** Rabbit Monoclonal Antibody

HostRabbitApplicationWB,ELISAReactivityHuman,Rat

### **Performance**

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type
Buffer	preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.
	Avoid freeze / thaw cycle.
Purification	Affinity purification

### **Immunogen**

Gene Name HAMP

Hepcidin; Liver-expressed antimicrobial peptide 1; LEAP-1; Putative liver tumor Alternative Names

regressor; PLTR; Hepcidin-25; Hepc25; Hepcidin-20; Hepc20; HAMP; HEPC; LEAP1;

 Gene ID
 57817.0

 SwissProt ID
 P81172.

# **Application**

**Dilution Ratio** WB 1:500-1:2000

Molecular Weight 9kDa

# Product Name: Hepcidin (16X4) Rabbit Monoclonal

**Antibody** 

Catalog #: AMRe11984




## **Background**

Hepcidin, also known as HAMP, HEPC, LEAP1 or HFE2B, is an 84 amino acid secreted protein that regulates iron-related signaling events. Acts by promoting endocytosis and degradation of ferroportin, leading to the retention of iron in iron-exporting cells and decreased flow of iron into plasma. Liver-produced hormone that constitutes the main circulating regulator of iron absorption and distribution across tissues. Acts by promoting endocytosis and degradation of ferroportin/SLC40A1, leading to the retention of iron in iron-exporting cells and decreased flow of iron into plasma (PubMed:<a href="http://www.uniprot.org/citations/22682227" target="\_blank">22682227</a>, PubMed:<a href="http://www.uniprot.org/citations/29237594" target="\_blank">29237594</a>, PubMed:<a href="http://www.uniprot.org/citations/32814342" target="\_blank">32814342</a>). Controls the major flows of iron into plasma: absorption of dietary iron in the intestine, recycling of iron by macrophages, which phagocytose old erythrocytes and other cells, and mobilization of stored iron from hepatocytes (PubMed:<a href="http://www.uniprot.org/citations/22306005" target=" blank">22306005</a>, a>).

### Research Area

### **Image Data**



Western blot analysis of Hepcidin expression in Human Hepcidin Full-length Recombinant Protein (GFP Tagged).

#### Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838