

# Summary

Production Name	Interferon gamma (16O15) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human,Mouse,Rat

#### Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

## Immunogen

Gene Name	IFNG
Alternative Names	IFG; IFI; IFN gamma; IFN, immune; IFN-gamma; IFNG; Immune interferon; Interferon
	gamma;
Gene ID	3458.0
SwissProt ID	P01579.

# Application

Dilution Ratio	WB 1:500-1:2000
Molecular Weight	19kDa



## Background

Interferon (IFN)-y is an antiviral and antiparasitic agent produced by CD4+/CD8+ lymphocytes and natural killer cells that undergo activation by antigens, mitogens or alloantigens. It is a potent activator of macrophages, it has antiproliferative effects on transformed cells and it can potentiate the antiviral and antitumor effects of the type I interferons. Type II interferon produced by immune cells such as T-cells and NK cells that plays crucial roles in antimicrobial, antiviral, and antitumor responses by activating effector immune cells and enhancing antigen presentation (PubMed:<a href="http://www.uniprot.org/citations/16914093" target=" blank">16914093</a>, PubMed:<a href="http://www.uniprot.org/citations/8666937" target=" blank">8666937</a>). Primarily signals through the JAK-STAT pathway after interaction with its receptor IFNGR1 to affect gene regulation (PubMed:<a href="http://www.uniprot.org/citations/8349687" target=" blank">8349687</a>). Upon IFNG binding, IFNGR1 intracellular domain opens out to allow association of downstream signaling components JAK2, JAK1 and STAT1, leading to STAT1 activation, nuclear translocation and transcription of IFNG-regulated genes. Many of the induced genes are transcription factors such as IRF1 that are able to further drive regulation of a next wave of transcription (PubMed:<a href="http://www.uniprot.org/citations/16914093" target=" blank">16914093</a>). Plays a role in class I antigen presentation pathway by inducing a replacement of catalytic proteasome subunits with immunoproteasome subunits (PubMed:<a href="http://www.uniprot.org/citations/8666937" target=" blank">8666937</a>). In turn, increases the quantity, quality, and repertoire of peptides for class I MHC loading (PubMed: <a href="http://www.uniprot.org/citations/8163024" target=" blank">8163024</a>). Increases the efficiency of peptide generation also by inducing the expression of activator PA28 that associates with the proteasome and alters its proteolytic cleavage preference (PubMed: <a href="http://www.uniprot.org/citations/11112687" target=" blank">11112687</a>). Upregulates as well MHC II complexes on the cell surface by promoting expression of several key molecules such as cathepsins B/CTSB, H/CTSH, and L/CTSL (PubMed: <a href="http://www.uniprot.org/citations/7729559" target=" blank">7729559</a>). Participates in the regulation of hematopoietic stem cells during development and under homeostatic conditions by affecting their development, quiescence, and differentiation (By similarity).

### **Research Area**

#### **Image Data**





Western blot analysis of Interferon gamma expression in Jurkat cell lysate.

#### Note

For research use only.