

Summary

lonal Antibody	
Rabbit Monoclonal Antibody	

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New typepreservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	JAG1
Alternative Names	JAG1; AGS; AHD; Alagille syndrome; CD339 antigen; HJ1; Jagged1; JAGL1; Jagged;
Alternative Names	Jagged 1; Protein jagged-1; AWS; CD339; Soluble protein jagged;
Gene ID	182.0
SwissProt ID	P78504.

Application

Dilution Ratio	WB 1:500-1:2000
Molecular Weight	134kDa




Background

Ligand for multiple Notch receptors and involved in the mediation of Notch signaling. May be involved in cell-fate decisions during hematopoiesis. Seems to be involved in early and late stages of mammalian cardiovascular development. Inhibits myoblast differentiation (By similarity). Enhances fibroblast growth factor-induced angiogenesis (in vitro). Ligand for multiple Notch receptors and involved in the mediation of Notch signaling (PubMed:18660822, PubMed:20437614). May be involved in cell-fate decisions during hematopoiesis (PubMed:20437614). May be involved in cell-fate decisions during hematopoiesis (PubMed:20437614). May be involved in cell-fate decisions during hematopoiesis (PubMed:20437614). May be involved in cell-fate decisions during hematopoiesis (PubMed:9462510). Seems to be involved in early and late stages of mammalian cardiovascular development. Inhibits myoblast differentiation (By similarity). Enhances fibroblast growth factor-induced angiogenesis (in vitro).

Research Area

Image Data



Western blot analysis of JAG1 expression in HepG2 cell lysate.

Note

For research use only.