

**Product Name: KDM5A (6A16) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe12971**



## Summary

|                        |   |
|------------------------|---|
| <b>Production Name</b> | KDM5A (6A16) Rabbit Monoclonal Antibody |
| <b>Description</b>     | Rabbit Monoclonal Antibody              |
| <b>Host</b>            | Rabbit                                  |
| <b>Application</b>     | WB                                      |
| <b>Reactivity</b>      | Human,Mouse                             |

## Performance

|                     |  |
|---------------------|--|
| <b>Conjugation</b>  | Unconjugated   |
| <b>Modification</b> | Unmodified   |
| <b>Isotype</b>      | IgG  |
| <b>Clonality</b>    | Monoclonal   |
| <b>Form</b>         | Liquid   |
| <b>Storage</b>      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.   |
| <b>Buffer</b>       | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |
| <b>Purification</b> | Affinity purification  |

## Immunogen

|                          |                              |
|--------------------------|------------------------------|
| <b>Gene Name</b>         | KDM5A                        |
| <b>Alternative Names</b> | JARID1A; Kdm5a; RBBP2; RBP2; |
| <b>Gene ID</b>           | 5927.0                       |
| <b>SwissProt ID</b>      | P29375.                      |

## Application

|                         |                 |
|-------------------------|-----------------|
| <b>Dilution Ratio</b>   | WB 1:500-1:2000 |
| <b>Molecular Weight</b> | 192kDa          |

**Product Name: KDM5A (6A16) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe12971**




## Background

Histone demethylase that specifically demethylates 'Lys-4' of histone H3, thereby playing a central role in histone code. Does not demethylate histone H3 'Lys-9', H3 'Lys-27', H3 'Lys-36', H3 'Lys-79' or H4 'Lys-20'. Demethylates trimethylated and dimethylated but not monomethylated H3 'Lys-4'. Histone demethylase that specifically demethylates 'Lys-4' of histone H3, thereby playing a central role in histone code. Does not demethylate histone H3 'Lys-9', H3 'Lys-27', H3 'Lys-36', H3 'Lys-79' or H4 'Lys-20'. Demethylates trimethylated and dimethylated but not monomethylated H3 'Lys-4'. Regulates specific gene transcription through DNA-binding on 5'-CCGCCC-3' motif (PubMed:<a href="http://www.uniprot.org/citations/18270511" target="\_blank">18270511</a>). May stimulate transcription mediated by nuclear receptors. Involved in transcriptional regulation of Hox proteins during cell differentiation (PubMed:<a href="http://www.uniprot.org/citations/19430464" target="\_blank">19430464</a>). May participate in transcriptional repression of cytokines such as CXCL12. Plays a role in the regulation of the circadian rhythm and in maintaining the normal periodicity of the circadian clock. In a histone demethylase-independent manner, acts as a coactivator of the CLOCK-ARNTL/BMAL1-mediated transcriptional activation of PER1/2 and other clock-controlled genes and increases histone acetylation at PER1/2 promoters by inhibiting the activity of HDAC1 (By similarity). Seems to act as a transcriptional corepressor for some genes such as MT1F and to favor the proliferation of cancer cells (PubMed:<a href="http://www.uniprot.org/citations/27427228" target="\_blank">27427228</a>).

## Research Area

## Image Data



Western blot analysis of KDM5A / Jarid1A / RBBP2 expression in (1) HEK293 cell lysate; (2) Mouse spleen lysate.

## Note

For research use only.