

Catalog #: AMRe13292



Summary

LGR5/GPR49 (2F16) Rabbit Monoclonal Antibody **Production Name**

Description Rabbit Monoclonal Antibody

Host Rabbit **Application** WB,ELISA

Reactivity Human, Mouse, Rat

Performance

| Conjugation | Unconjugated |
|--------------|------------------------------------------------------------------------------------------|
| Modification | Unmodified |
| Isotype | IgG |
| Clonality | Monoclonal |
| Form | Liquid |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |
| | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type |
| Buffer | preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. |
| | Avoid freeze / thaw cycle. |
| Purification | Affinity purification |

Immunogen

Gene Name LGR5

Alternative Names FEX; GPR49; GPR67; GRP49; LGR5; HG38;8

Gene ID 8549.0 SwissProt ID O75473.

Application

Dilution Ratio WB 1:500-1:2000

Molecular Weight 100kDa

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: LGR5/GPR49 (2F16) Rabbit Monoclonal

Antibody

Catalog #: AMRe13292




Background

Receptor for R-spondins that potentiates the canonical Wnt signaling pathway and acts as a stem cell marker of the intestinal epithelium and the hair follicle. Upon binding to R-spondins (RSPO1, RSPO2, RSPO3 or RSPO4), associates with phosphorylated LRP6 and frizzled receptors that are activated by extracellular Wnt receptors, triggering the canonical Wnt signaling pathway to increase expression of target genes. Receptor for R-spondins that potentiates the canonical Wnt signaling pathway and acts as a stem cell marker of the intestinal epithelium and the hair follicle. Upon binding to R-spondins (RSPO1, RSPO2, RSPO3 or RSPO4), associates with phosphorylated LRP6 and frizzled receptors that are activated by extracellular Wnt receptors, triggering the canonical Wnt signaling pathway to increase expression of target genes. In contrast to classical G-protein coupled receptors, does not activate heterotrimeric G-proteins to transduce the signal. Involved in the development and/or maintenance of the adult intestinal stem cells during postembryonic development.

Research Area

Image Data



Western blot analysis of GPR49 expression in Human fetal skeletal muscle lysate.

Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838