

Product Name: MALT1 (7A3) Rabbit Monoclonal Antibody
Catalog #: AMRe13606



Summary

Production Name	MALT1 (7A3) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	MALT1 {ECO:0000303 PubMed:10523859, ECO:0000312 HGNC:HGNC:6819}
Alternative Names	Malt1; MLT1; Mucosa associated lymphoid tissue lymphoma translocation gene 1; Paracaspase;
Gene ID	10892.0
SwissProt ID	Q9UDY8.

Application

Dilution Ratio	WB 1:500-1:2000
Molecular Weight	92kDa

Product Name: MALT1 (7A3) Rabbit Monoclonal Antibody
Catalog #: AMRe13606




Background

Enhances BCL10-induced activation of NF-kappa-B. Involved in nuclear export of BCL10. Binds to TRAF6, inducing TRAF6 oligomerization and activation of its ligase activity. Has ubiquitin ligase activity. MALT1-dependent BCL10 cleavage plays an important role in T-cell antigen receptor-induced integrin adhesion. Protease that enhances BCL10-induced activation: acts via formation of CBM complexes that channel adaptive and innate immune signaling downstream of CARD domain-containing proteins (CARD9, CARD11 and CARD14) to activate NF-kappa-B and MAP kinase p38 pathways which stimulate expression of genes encoding pro-inflammatory cytokines and chemokines (PubMed: [11262391](http://www.uniprot.org/citations/11262391), PubMed: [18264101](http://www.uniprot.org/citations/18264101), PubMed: [24074955](http://www.uniprot.org/citations/24074955)). Mediates BCL10 cleavage: MALT1-dependent BCL10 cleavage plays an important role in T-cell antigen receptor-induced integrin adhesion (PubMed: [11262391](http://www.uniprot.org/citations/11262391), PubMed: [18264101](http://www.uniprot.org/citations/18264101)). Involved in the induction of T helper 17 cells (Th17) differentiation (PubMed: [11262391](http://www.uniprot.org/citations/11262391), PubMed: [18264101](http://www.uniprot.org/citations/18264101)). Cleaves RC3H1 and ZC3H12A in response to T-cell receptor (TCR) stimulation which releases their cooperatively repressed targets to promote Th17 cell differentiation (By similarity). Also mediates cleavage of N4BP1 in T-cells following TCR-mediated activation, leading to N4BP1 inactivation (PubMed: [31133753](http://www.uniprot.org/citations/31133753)). May also have ubiquitin ligase activity: binds to TRAF6, inducing TRAF6 oligomerization and activation of its ligase activity (PubMed: [14695475](http://www.uniprot.org/citations/14695475)).

Research Area

Image Data



**Product Name: MALT1 (7A3) Rabbit Monoclonal
Antibody**
Catalog #: AMRe13606



Western blot analysis of MALT1 expression in Jurkat cell lysate.

Note

For research use only.