

# Summary

Production Name	Mov10 (4D14) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human, Mouse, Rat

#### Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New typepreservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.Avoid freeze / thaw cycle.
Purification	Affinity purification

# Immunogen

Gene Name	MOV10
Alternative Names	gb110; MOV 10;
Gene ID	4343.0
SwissProt ID	Q9HCE1.

# Application

Dilution Ratio	WB 1:500-1:2000
Molecular Weight	114kDa




### Background

MOV10 may be an helicase with an important function in development and/or control of cell proliferation. RNA silencing processes are guided by small RNAs known as siRNAs and microRNAs (miRNAs). They reside in ribonucleoprotein complexes, which guide the cleavage of complementary mRNAs or affect stability and translation of partial complementary mRNAs. 5' to 3' RNA helicase contributing to UPF1 mRNA target degradation by translocation along 3' UTRs (PubMed:<a href="http://www.uniprot.org/citations/24726324" target="\_blank">24726324</a>). Required for microRNA (miRNA)mediated gene silencing by the RNA-induced silencing complex (RISC). Required for both miRNA-mediated translational repression and miRNA-mediated cleavage of complementary mRNAs by RISC (PubMed: <a href="http://www.uniprot.org/citations/16289642" target=" blank">16289642</a>, PubMed:<a href="http://www.uniprot.org/citations/17507929" target=" blank">17507929</a>, PubMed:<a href="http://www.uniprot.org/citations/22791714" target=" blank">22791714</a>). In cooperation with FMR1, regulates miRNA-mediated translational repression by AGO2 (PubMed: <a href="http://www.uniprot.org/citations/25464849" target=" blank">25464849</a>). Restricts retrotransposition of long interspersed element-1 (LINE-1) in cooperation with TUT4 and TUT7 counteracting the RNA chaperonne activity of L1RE1 (PubMed:<a href="http://www.uniprot.org/citations/30122351" target=" blank">30122351</a>, PubMed:<a href="http://www.uniprot.org/citations/23093941" target=" blank">23093941</a>). Facilitates LINE-1 uridylation by TUT4 and TUT7 (PubMed: <a href="http://www.uniprot.org/citations/30122351" target=" blank">30122351</a>). Required for embryonic viability and for normal central nervous system development and function. Plays two critical roles in early brain development: suppresses retroelements in the nucleus by directly inhibiting cDNA synthesis, while regulates cytoskeletal mRNAs to influence neurite outgrowth in the cytosol (By similarity). May function as a messenger ribonucleoprotein (mRNP) clearance factor (PubMed:<a href="http://www.uniprot.org/citations/24726324" target=" blank">24726324</a>). Exhibits antiviral activity against dengue virus (DENV) (PubMed: <a href="http://www.uniprot.org/citations/27974568" target=" blank">27974568</a>).

## **Research Area**

#### **Image Data**





Western blot analysis of Mov10 expression in (1) 293 cell lysate; (2) NIH/3T3 cell lysate.

#### Note

For research use only.