

Summary

Production Name	OGG1 (15C8) Rabbit Monoclonal Antibody	
Description	Rabbit Monoclonal Antibody	
Host	Rabbit	
Application	WB,ELISA	
Reactivity	Human	

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New typepreservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	OGG1
Alternative Names	8 hydroxyguanine DNA glycosylase; AP lyase; HMMH; HOGG1; MMH; MUTM; N
	glycosylase; Ogg1; OGH 1;
Gene ID	4968.0
SwissProt ID	O15527.

Application

Dilution Ratio	WB 1:500-1:2000
Molecular Weight	39kDa




Background

DNA repair enzyme that incises DNA at 8-oxoG residues. Excises 7,8-dihydro-8-oxoguanine and 2,6-diamino-4-hydroxy-5-N-methylformamidopyrimidine (FAPY) from damaged DNA. Has a beta-lyase activity that nicks DNA 3' to the lesion. DNA repair enzyme that incises DNA at 8-oxoG residues. Excises 7,8-dihydro-8-oxoguanine and 2,6-diamino-4-hydroxy-5-Nmethylformamidopyrimidine (FAPY) from damaged DNA. Has a beta-lyase activity that nicks DNA 3' to the lesion.

Research Area

Image Data



Western blot analysis of OGG1 expression in HepG2 cell lysate.

Note

For research use only.