

**Product Name: P70 S6 Kinase alpha (10D3) Rabbit  
Monoclonal Antibody  
Catalog #: AMRe15663**



## Summary

<b>Production Name</b>	P70 S6 Kinase alpha (10D3) Rabbit Monoclonal Antibody
<b>Description</b>	Rabbit Monoclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,ELISA
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	RPS6KB1
<b>Alternative Names</b>	p70 alpha; P70 beta 1; p70 S6K; P70S6K; P70S6K1; p70S6Kb; PS6K; RPS6KB1; S6K; S6K1; STK14A;
<b>Gene ID</b>	6198.0
<b>SwissProt ID</b>	P23443.

## Application

<b>Dilution Ratio</b>	WB 1:500-1:1000
<b>Molecular Weight</b>	59kDa

**Product Name: P70 S6 Kinase alpha (10D3) Rabbit  
Monoclonal Antibody  
Catalog #: AMRe15663**



## Background

Acts to integrate nutrient and growth factor signals in regulation of protein synthesis, cell proliferation, cell growth, cell cycle progression and cell survival. Downstream effector of the mTOR signaling pathway. Serine/threonine-protein kinase that acts downstream of mTOR signaling in response to growth factors and nutrients to promote cell proliferation, cell growth and cell cycle progression. Regulates protein synthesis through phosphorylation of EIF4B, RPS6 and EEF2K, and contributes to cell survival by repressing the pro-apoptotic function of BAD. Under conditions of nutrient depletion, the inactive form associates with the EIF3 translation initiation complex. Upon mitogenic stimulation, phosphorylation by the mammalian target of rapamycin complex 1 (mTORC1) leads to dissociation from the EIF3 complex and activation. The active form then phosphorylates and activates several substrates in the pre-initiation complex, including the EIF2B complex and the cap-binding complex component EIF4B. Also controls translation initiation by phosphorylating a negative regulator of EIF4A, PDCD4, targeting it for ubiquitination and subsequent proteolysis. Promotes initiation of the pioneer round of protein synthesis by phosphorylating POLDIP3/SKAR. In response to IGF1, activates translation elongation by phosphorylating EEF2 kinase (EEF2K), which leads to its inhibition and thus activation of EEF2. Also plays a role in feedback regulation of mTORC2 by mTORC1 by phosphorylating RICTOR, resulting in the inhibition of mTORC2 and AKT1 signaling. Mediates cell survival by phosphorylating the pro-apoptotic protein BAD and suppressing its pro-apoptotic function. Phosphorylates mitochondrial URI1 leading to dissociation of a URI1-PPP1CC complex. The free mitochondrial PPP1CC can then dephosphorylate RPS6KB1 at Thr-412, which is proposed to be a negative feedback mechanism for the RPS6KB1 anti-apoptotic function. Mediates TNF-alpha-induced insulin resistance by phosphorylating IRS1 at multiple serine residues, resulting in accelerated degradation of IRS1. In cells lacking functional TSC1-2 complex, constitutively phosphorylates and inhibits GSK3B. May be involved in cytoskeletal rearrangement through binding to neurabin. Phosphorylates and activates the pyrimidine biosynthesis enzyme CAD, downstream of MTOR (PubMed:<a

href="http://www.uniprot.org/citations/11500364" target="\_blank">11500364</a>, PubMed:<a href="http://www.uniprot.org/citations/12801526" target="\_blank">12801526</a>, PubMed:<a href="http://www.uniprot.org/citations/14673156" target="\_blank">14673156</a>, PubMed:<a href="http://www.uniprot.org/citations/15071500" target="\_blank">15071500</a>, PubMed:<a href="http://www.uniprot.org/citations/15341740" target="\_blank">15341740</a>, PubMed:<a href="http://www.uniprot.org/citations/16286006" target="\_blank">16286006</a>, PubMed:<a href="http://www.uniprot.org/citations/17052453" target="\_blank">17052453</a>, PubMed:<a href="http://www.uniprot.org/citations/17053147" target="\_blank">17053147</a>, PubMed:<a href="http://www.uniprot.org/citations/17936702" target="\_blank">17936702</a>, PubMed:<a href="http://www.uniprot.org/citations/18952604" target="\_blank">18952604</a>, PubMed:<a href="http://www.uniprot.org/citations/19085255" target="\_blank">19085255</a>, PubMed:<a href="http://www.uniprot.org/citations/19720745" target="\_blank">19720745</a>, PubMed:<a href="http://www.uniprot.org/citations/19935711" target="\_blank">19935711</a>, PubMed:<a


**Product Name: P70 S6 Kinase alpha (10D3) Rabbit  
Monoclonal Antibody  
Catalog #: AMRe15663**

---

<http://www.uniprot.org/citations/19995915> target="\_blank">19995915</a>, PubMed:<a href="http://www.uniprot.org/citations/23429703" target="\_blank">23429703</a>). Following activation by mTORC1, phosphorylates EPRS and thereby plays a key role in fatty acid uptake by adipocytes and also most probably in interferon-gamma-induced translation inhibition (PubMed:<a href="http://www.uniprot.org/citations/28178239" target="\_blank">28178239</a>).

## Research Area

## Image Data



Western blot analysis of P70 S6 Kinase alpha expression in (1) HeLa cell lysate; (2) Raw264.7 cell lysate; (3) PC12 cell lysate.

## Note

For research use only.