

Product Name: PABPN1 (9U19) Rabbit Monoclonal Antibody
Catalog #: AMRe15683



Summary

Production Name	PABPN1 (9U19) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
Purification	Affinity purification

Immunogen

Gene Name	PABPN1
Alternative Names	Nuclear poly(A)-binding protein 1; PAB2; PABII; PABP2; PABPII; Pabpn1; poly(A) binding protein nuclear 1;
Gene ID	8106.0
SwissProt ID	Q86U42.A synthetic peptide of human PABPN1

Application

Dilution Ratio	WB: 1:1000-1:5000
Molecular Weight	33kDa

Product Name: PABPN1 (9U19) Rabbit Monoclonal Antibody
Catalog #: AMRe15683




Background

Involved in the 3'-end formation of mRNA precursors (pre-mRNA) by the addition of a poly(A) tail of 200-250 nt to the upstream cleavage product. Stimulates poly(A) polymerase (PAPOLA) conferring processivity on the poly(A) tail elongation reaction and controls also the poly(A) tail length. Increases the affinity of poly(A) polymerase for RNA. Involved in the 3'-end formation of mRNA precursors (pre- mRNA) by the addition of a poly(A) tail of 200-250 nt to the upstream cleavage product (By similarity). Stimulates poly(A) polymerase (PAPOLA) conferring processivity on the poly(A) tail elongation reaction and controls also the poly(A) tail length (By similarity). Increases the affinity of poly(A) polymerase for RNA (By similarity). Is also present at various stages of mRNA metabolism including nucleocytoplasmic trafficking and nonsense-mediated decay (NMD) of mRNA. Cooperates with SKIP to synergistically activate E-box-mediated transcription through MYOD1 and may regulate the expression of muscle- specific genes (PubMed:11371506). Binds to poly(A) and to poly(G) with high affinity (By similarity). May protect the poly(A) tail from degradation (By similarity). Subunit of the trimeric poly(A) tail exosome targeting (PAXT) complex, a complex that directs a subset of long and polyadenylated poly(A) RNAs for exosomal degradation. The RNA exosome is fundamental for the degradation of RNA in eukaryotic nuclei. Substrate targeting is facilitated by its cofactor MTREX, which links to RNA-binding protein adapters (PubMed:27871484).

Research Area

Image Data



Western blot detection of PABPN1 in Jurkat, HeLa, COS7, Raji cell lysates using PABPN1 antibody (1:1000 diluted).

Note

For research use only.