

Product Name: PARK7 (9L2) Rabbit Monoclonal Antibody
Catalog #: AMRe15756

Summary

Production Name	PARK7 (9L2) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
Purification	Affinity purification

Immunogen

Gene Name	PARK7
Alternative Names	Protein DJ-1; SP22; Protein DJ-1; Oncogene DJ1; Parkinson disease protein 7; PARK7;
Gene ID	11315.0
SwissProt ID	Q99497.A synthetic peptide of human PARK7/DJ1

Application

Dilution Ratio	WB: 1:2000-1:10000
Molecular Weight	20kDa

Background

Product Name: PARK7 (9L2) Rabbit Monoclonal Antibody
Catalog #: AMRe15756



Plays a role in regulating expression or stability of the mitochondrial uncoupling proteins SLC25A14 and SLC25A27 in dopaminergic neurons of the substantia nigra pars compacta and attenuates the oxidative stress induced by calcium entry into the neurons via L-type channels during pacemaking. It cooperates with Ras to increase cell transformation, it positively regulates transcription of the androgen receptor, and it may function as an indicator of oxidative stress. Multifunctional protein with controversial molecular function which plays an important role in cell protection against oxidative stress and cell death acting as oxidative stress sensor and redox- sensitive chaperone and protease (PubMed:[17015834](http://www.uniprot.org/citations/17015834), PubMed:[20304780](http://www.uniprot.org/citations/20304780), PubMed:[18711745](http://www.uniprot.org/citations/18711745), PubMed:[12796482](http://www.uniprot.org/citations/12796482), PubMed:[19229105](http://www.uniprot.org/citations/19229105), PubMed:[25416785](http://www.uniprot.org/citations/25416785), PubMed:[26995087](http://www.uniprot.org/citations/26995087), PubMed:[28993701](http://www.uniprot.org/citations/28993701)). It is involved in neuroprotective mechanisms like the stabilization of NFE2L2 and PINK1 proteins, male fertility as a positive regulator of androgen signaling pathway as well as cell growth and transformation through, for instance, the modulation of NF-kappa-B signaling pathway (PubMed:[12612053](http://www.uniprot.org/citations/12612053), PubMed:[15502874](http://www.uniprot.org/citations/15502874), PubMed:[14749723](http://www.uniprot.org/citations/14749723), PubMed:[17015834](http://www.uniprot.org/citations/17015834), PubMed:[21097510](http://www.uniprot.org/citations/21097510), PubMed:[18711745](http://www.uniprot.org/citations/18711745)). Has been described as a protein and nucleotide deglycase that catalyzes the deglycation of the Maillard adducts formed between amino groups of proteins or nucleotides and reactive carbonyl groups of glyoxals (PubMed:[25416785](http://www.uniprot.org/citations/25416785), PubMed:[28596309](http://www.uniprot.org/citations/28596309)). But this function is rebuted by other works (PubMed:[27903648](http://www.uniprot.org/citations/27903648), PubMed:[31653696](http://www.uniprot.org/citations/31653696)). As a protein deglycase, repairs methylglyoxal- and glyoxal-glycated proteins, and releases repaired proteins and lactate or glycolate, respectively. Deglycates cysteine, arginine and lysine residues in proteins, and thus reactivates these proteins by reversing glycation by glyoxals. Acts on early glycation intermediates (hemithioacetals and aminocarbinols), preventing the formation of advanced glycation endproducts (AGE) that cause irreversible damage (PubMed:[25416785](http://www.uniprot.org/citations/25416785), PubMed:[28013050](http://www.uniprot.org/citations/28013050), PubMed:[26995087](http://www.uniprot.org/citations/26995087)). Also functions as a nucleotide

Product Name: PARK7 (9L2) Rabbit Monoclonal Antibody
Catalog #: AMRe15756


deglycase able to repair glycated guanine in the free nucleotide pool (GTP, GDP, GMP, dGTP) and in DNA and RNA. Is thus involved in a major nucleotide repair system named guanine glycation repair (GG repair), dedicated to reversing methylglyoxal and glyoxal damage via nucleotide sanitization and direct nucleic acid repair (PubMed: [28596309](http://www.uniprot.org/citations/28596309)). Protects histones from adduction by methylglyoxal, controls the levels of methylglyoxal- derived argininine modifications on chromatin (PubMed: [30150385](http://www.uniprot.org/citations/30150385)). Able to remove the glycations and restore histone 3, histone glycation disrupts both local and global chromatin architecture by altering histone-DNA interactions as well as histone acetylation and ubiquitination levels (PubMed: [30150385](http://www.uniprot.org/citations/30150385)), PubMed: [30894531](http://www.uniprot.org/citations/30894531)). Displays a very low glyoxalase activity that may reflect its deglycase activity (PubMed: [22523093](http://www.uniprot.org/citations/22523093)), PubMed: [31653696](http://www.uniprot.org/citations/31653696), PubMed: [28993701](http://www.uniprot.org/citations/28993701)). Eliminates hydrogen peroxide and protects cells against hydrogen peroxide-induced cell death (PubMed: [16390825](http://www.uniprot.org/citations/16390825)). Required for correct mitochondrial morphology and function as well as for autophagy of dysfunctional mitochondria (PubMed: [19229105](http://www.uniprot.org/citations/19229105)), PubMed: [16632486](http://www.uniprot.org/citations/16632486)). Plays a role in regulating expression or stability of the mitochondrial uncoupling proteins SLC25A14 and SLC25A27 in dopaminergic neurons of the substantia nigra pars compacta and attenuates the oxidative stress induced by calcium entry into the neurons via L-type channels during pacemaking (PubMed: [18711745](http://www.uniprot.org/citations/18711745)). Regulates astrocyte inflammatory responses, may modulate lipid rafts-dependent endocytosis in astrocytes and neuronal cells (PubMed: [23847046](http://www.uniprot.org/citations/23847046)). In pancreatic islets, involved in the maintenance of mitochondrial reactive oxygen species (ROS) levels and glucose homeostasis in an age- and diet dependent manner. Protects pancreatic beta cells from cell death induced by inflammatory and cytotoxic setting (By similarity). Binds to a number of mRNAs containing multiple copies of GG or CC motifs and partially inhibits their translation but dissociates following oxidative stress (PubMed: [18626009](http://www.uniprot.org/citations/18626009)). Metal-binding protein able to bind copper as well as toxic mercury ions, enhances the cell protection mechanism against induced metal toxicity (PubMed: [23792957](http://www.uniprot.org/citations/23792957)). In macrophages, interacts with the NADPH oxidase subunit NCF1 to direct NADPH oxidase-dependent ROS production, and protects against sepsis (By similarity).

Research Area

Product Name: PARK7 (9L2) Rabbit Monoclonal Antibody
Catalog #: AMRe15756



Image Data



Western blot detection of PARK7/DJ1 in K562,Rat Brain,C6,3T3,HeLa cell lysates using PARK7/DJ1 antibody(1:1000 diluted).

Note

For research use only.