

**Product Name: PEF1 (18D15) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe15954**

---

## Summary

<b>Production Name</b>	PEF1 (18D15) Rabbit Monoclonal Antibody
<b>Description</b>	Rabbit Monoclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	PEF1
<b>Alternative Names</b>	ABP32; PEF; pef1; PEF1A; Peflin;
<b>Gene ID</b>	553115.0
<b>SwissProt ID</b>	Q9UBV8.A synthetic peptide of human PEF1

## Application

<b>Dilution Ratio</b>	WB: 1:1000-1:5000
<b>Molecular Weight</b>	30kDa

## Background

---


**Product Name: PEF1 (18D15) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe15954**

---

Calcium-binding protein that acts as an adapter that bridges unrelated proteins or stabilizes weak protein-protein complexes in response to calcium. Together with PDCD6, acts as calcium-dependent adapter for the BCR(KLHL12) complex, a complex involved in endoplasmic reticulum (ER)-Golgi transport by regulating the size of COPII coats (PubMed:27716508). Calcium-binding protein that acts as an adapter that bridges unrelated proteins or stabilizes weak protein-protein complexes in response to calcium. Together with PDCD6, acts as calcium-dependent adapter for the BCR(KLHL12) complex, a complex involved in endoplasmic reticulum (ER)-Golgi transport by regulating the size of COPII coats (PubMed:<a href="http://www.uniprot.org/citations/27716508" target="\_blank">27716508</a>). In response to cytosolic calcium increase, the heterodimer formed with PDCD6 interacts with, and bridges together the BCR(KLHL12) complex and SEC31 (SEC31A or SEC31B), promoting monoubiquitination of SEC31 and subsequent collagen export, which is required for neural crest specification (PubMed:<a href="http://www.uniprot.org/citations/27716508" target="\_blank">27716508</a>). Its role in the heterodimer formed with PDCD6 is however unclear: some evidence shows that PEF1 and PDCD6 work together and promote association between PDCD6 and SEC31 in presence of calcium (PubMed:<a href="http://www.uniprot.org/citations/27716508" target="\_blank">27716508</a>). Other reports show that PEF1 dissociates from PDCD6 in presence of calcium, and may act as a negative regulator of PDCD6 (PubMed:<a href="http://www.uniprot.org/citations/11278427" target="\_blank">11278427</a>). Also acts as a negative regulator of ER-Golgi transport; possibly by inhibiting interaction between PDCD6 and SEC31 (By similarity).

## Research Area

## Image Data



Western blot detection of PEF1 in K562,Rat Brain,C6,3T3,HeLa cell lysates using PEF1 antibody(1:1000 diluted).

## Note

For research use only.