

**Product Name: PI 3 Kinase p85 alpha (17X10) Rabbit  
Monoclonal Antibody  
Catalog #: AMRe16094**



## Summary

<b>Production Name</b>	PI 3 Kinase p85 alpha (17X10) Rabbit Monoclonal Antibody
<b>Description</b>	Rabbit Monoclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	PIK3R1
<b>Alternative Names</b>	GRB1; P85A; PI3-kinase p85-alpha; PI3K; PI3K p85-alpha; ptdIns-3-kinase p85-alpha;
<b>Gene ID</b>	5295.0
<b>SwissProt ID</b>	P27986.A synthetic peptide of human PI 3 Kinase p85 alpha

## Application

<b>Dilution Ratio</b>	WB: 1:1000
<b>Molecular Weight</b>	84kDa

## Background


**Product Name: PI 3 Kinase p85 alpha (17X10) Rabbit  
Monoclonal Antibody  
Catalog #: AMRe16094**



Binds to activated (phosphorylated) protein-Tyr kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Necessary for the insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues. Plays an important role in signaling in response to FGFR1, FGFR2, FGFR3, FGFR4, KITLG/SCF, KIT, PDGFRA and PDGFRB. Likewise, plays a role in ITGB2 signaling. Binds to activated (phosphorylated) protein-Tyr kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Necessary for the insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues. Plays an important role in signaling in response to FGFR1, FGFR2, FGFR3, FGFR4, KITLG/SCF, KIT, PDGFRA and PDGFRB. Likewise, plays a role in ITGB2 signaling (PubMed:<a href="http://www.uniprot.org/citations/17626883" target="\_blank">17626883</a>, PubMed:<a href="http://www.uniprot.org/citations/19805105" target="\_blank">19805105</a>, PubMed:<a href="http://www.uniprot.org/citations/7518429" target="\_blank">7518429</a>). Modulates the cellular response to ER stress by promoting nuclear translocation of XBP1 isoform 2 in a ER stress- and/or insulin-dependent manner during metabolic overloading in the liver and hence plays a role in glucose tolerance improvement (PubMed:<a href="http://www.uniprot.org/citations/20348923" target="\_blank">20348923</a>).

## Research Area

## Image Data



Western blot detection of PI 3 Kinase p85 alpha in HL-60,U2OS,C6 cell lysates using PI 3 Kinase p85 alpha antibody(1:1000 diluted).

## Note

For research use only.