

# Summary

Production Name	Podoplanin (11C17) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human, Mouse, Rat

#### Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.
	Avoid freeze / thaw cycle.
Purification	Affinity purification

### Immunogen

Gene Name	PDPN {ECO:0000312 EMBL:AAH14668.2}	
Alternative Names	Aggrus; Glycoprotein 36 KD; GP36; GP38; GP40; HT1A1; hT1alpha1; hT1alpha2; OTS8;	
	PA2.26; Pdpn; Podoplanin; T1 alpha; TI1A; TIA2;	
Gene ID	10630.0	
SwissProt ID	Q86YL7.	

# Application

Dilution Ratio	WB 1:500-1:2000
Molecular Weight	17kDa



## Background

May be involved in cell migration and/or actin cytoskeleton organization. When expressed in keratinocytes, induces changes in cell morphology with transfected cells showing an elongated shape, numerous membrane protrusions, major reorganization of the actin cytoskeleton, increased motility and decreased cell adhesion. Required for normal lung cell proliferation and alveolus formation at birth. Induces platelet aggregation. Mediates effects on cell migration and adhesion through its different partners. During development plays a role in blood and lymphatic vessels separation by binding CLEC1B, triggering CLEC1B activation in platelets and leading to platelet activation and/or aggregation (PubMed: <a href="http://www.uniprot.org/citations/14522983" target=" blank">14522983</a>, PubMed:<a href="http://www.uniprot.org/citations/15231832" target=" blank">15231832</a>, PubMed:<a href="http://www.uniprot.org/citations/17616532" target=" blank">17616532</a>, PubMed:<a href="http://www.uniprot.org/citations/18215137" target=" blank">18215137</a>, PubMed:<a href="http://www.uniprot.org/citations/17222411" target=" blank">17222411</a>). Interaction with CD9, on the contrary, attenuates platelet aggregation induced by PDPN (PubMed: <a href="http://www.uniprot.org/citations/18541721" target=" blank">18541721</a>). Through MSN or EZR interaction promotes epithelial- mesenchymal transition (EMT) leading to ERZ phosphorylation and triggering RHOA activation leading to cell migration increase and invasiveness (PubMed:<a href="http://www.uniprot.org/citations/17046996" target=" blank">17046996</a>, PubMed:<a href="http://www.uniprot.org/citations/21376833" target=" blank">21376833</a>). Interaction with CD44 promotes directional cell migration in epithelial and tumor cells (PubMed: <a href="http://www.uniprot.org/citations/20962267" target=" blank">20962267</a>). In lymph nodes (LNs), controls fibroblastic reticular cells (FRCs) adhesion to the extracellular matrix (ECM) and contraction of the actomyosin by maintaining ERM proteins (EZR; MSN and RDX) and MYL9 activation through association with unknown transmembrane proteins. Engagement of CLEC1B by PDPN promotes FRCs relaxation by blocking lateral membrane interactions leading to reduction of ERM proteins (EZR; MSN and RDX) and MYL9 activation (By similarity). Through binding with LGALS8 may participate in connection of the lymphatic endothelium to the surrounding extracellular matrix (PubMed: <a href="http://www.uniprot.org/citations/19268462" target=" blank">19268462</a>). In keratinocytes, induces changes in cell morphology showing an elongated shape, numerous membrane protrusions, major reorganization of the actin cytoskeleton, increased motility and decreased cell adhesion (PubMed:<a href="http://www.uniprot.org/citations/15515019" target=" blank">15515019</a>). Controls invadopodia stability and maturation leading to efficient degradation of the extracellular matrix (ECM) in tumor cells through modulation of RHOC activity in order to activate ROCK1/ROCK2 and LIMK1/LIMK2 and inactivation of CFL1 (PubMed: <a href="http://www.uniprot.org/citations/25486435" target=" blank">25486435</a>). Required for normal lung cell proliferation and alveolus formation at birth (By similarity). Does not function as a water channel or as a regulator of aquaporin-type water channels (PubMed:<a href="http://www.uniprot.org/citations/9651190" target=" blank">9651190</a>). Does not have any effect on folic acid or amino acid transport (By similarity).

Product Name: Podoplanin (11C17) Rabbit Monoclonal **Control Control Co** 

## **Research Area**

#### Image Data



Western blot analysis of Podoplanin expression in human placenta lysate.

#### Note

For research use only.