**Product Name: Smad2 (3F7) Rabbit Monoclonal** 

**Antibody** 

Catalog #: AMRe17990



# **Summary**

**Production Name** Smad2 (3F7) Rabbit Monoclonal Antibody

**Description** Rabbit Monoclonal Antibody

Host Rabbit
Application WB

**Reactivity** Human,Rat

### **Performance**

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
Purification	Affinity purification

### **Immunogen**

Gene Name SMAD2

JV18-1; MADH2; MADR2; Mad-related protein 2; Mothers against DPP homolog 2; Alternative Names

Mothers against decapentaplegic homolog 2; Smad2;

**Gene ID** 4087.0

**SwissProt ID** Q15796.A synthetic peptide of human Smad2

**Application** 

Dilution Ratio WB: 1:1000-1:5000

Molecular Weight 52kDa

 **Product Name: Smad2 (3F7) Rabbit Monoclonal** 

**Antibody** 

Catalog #: AMRe17990




### **Background**

The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. Receptor-regulated SMAD (R-SMAD) that is an intracellular signal transducer and transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinases. Binds the TRE element in the promoter region of many genes that are regulated by TGF-beta and, on formation of the SMAD2/SMAD4 complex, activates transcription. May act as a tumor suppressor in colorectal carcinoma. Positively regulates PDPK1 kinase activity by stimulating its dissociation from the 14-3-3 protein YWHAQ which acts as a negative regulator.

#### Research Area

## **Image Data**



Western blot detection of Smad2 in Hela, CHO-K1, Jurkat cell lysates using Smad2 antibody (1:500 diluted).

#### Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838