

**Product Name: Wilms Tumor Protein (13C17) Rabbit
Monoclonal Antibody
Catalog #: AMRe19901**



Summary

Production Name	Wilms Tumor Protein (13C17) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	WT1
Alternative Names	GUD; AWT1; WAGR; WT33; NPHS4; WIT-2; EWS-WT1;
Gene ID	7490.0
SwissProt ID	P19544.

Application

Dilution Ratio	WB 1:500-1:1000
Molecular Weight	49kDa

**Product Name: Wilms Tumor Protein (13C17) Rabbit
Monoclonal Antibody
Catalog #: AMRe19901**




Background

Has an essential role in the normal development of the urogenital system, and it is mutated in a small subset of patients with Wilm's tumors. This gene exhibits complex tissue-specific and polymorphic imprinting pattern, with biallelic, and monoallelic expression from the maternal and paternal alleles in different tissues. Multiple transcript variants have been described. In several variants, there is evidence for the use of a non-AUG (CUG) translation initiation site upstream of and in-frame with the first AUG. Transcription factor that plays an important role in cellular development and cell survival (PubMed:7862533). Recognizes and binds to the DNA sequence 5'-GCG(T/G)GGGCG-3' (PubMed:7862533, PubMed:17716689, PubMed:25258363). Regulates the expression of numerous target genes, including EPO. Plays an essential role for development of the urogenital system. It has a tumor suppressor as well as an oncogenic role in tumor formation. Function may be isoform-specific: isoforms lacking the KTS motif may act as transcription factors (PubMed:15520190). Isoforms containing the KTS motif may bind mRNA and play a role in mRNA metabolism or splicing (PubMed:16934801). Isoform 1 has lower affinity for DNA, and can bind RNA (PubMed:19123921).

Research Area

Image Data



Western blot analysis of WT1 expression in K562 cell lysate.

Note

For research use only.

**Product Name: Wilms Tumor Protein (13C17) Rabbit
Monoclonal Antibody
Catalog #: AMRe19901**

